



# Multi-dimensional Burgers equation with unbounded initial data: well-posedness and dispersive estimates

Denis Serre, Luis Silvestre

## ► To cite this version:

Denis Serre, Luis Silvestre. Multi-dimensional Burgers equation with unbounded initial data: well-posedness and dispersive estimates. 2018. ensl-01858016

**HAL Id: ensl-01858016**

**<https://ens-lyon.hal.science/ensl-01858016>**

Preprint submitted on 18 Aug 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Multi-dimensional Burgers equation with unbounded initial data: well-posedness and dispersive estimates

Denis Serre<sup>\*</sup>  
Luis Silvestre<sup>†‡</sup>

## Abstract

The Cauchy problem for a scalar conservation laws admits a unique entropy solution when the data  $u_0$  is a bounded measurable function (Kruzhkov). The semi-group  $(S_t)_{t \geq 0}$  is contracting in the  $L^1$ -distance.

For the multi-dimensional Burgers equation, we show that  $(S_t)_{t \geq 0}$  extends uniquely as a continuous semi-group over  $L^p(\mathbb{R}^n)$  whenever  $1 \leq p < \infty$ , and  $u(t) := S_t u_0$  is actually an entropy solution to the Cauchy problem. When  $p \leq q \leq \infty$  and  $t > 0$ ,  $S_t$  actually maps  $L^p(\mathbb{R}^n)$  into  $L^q(\mathbb{R}^n)$ .

These results are based upon new dispersive estimates. The ingredients are on the one hand Compensated Integrability, and on the other hand a De Giorgi-type iteration.

**Key words:** Dispersive estimates, Compensated integrability, Scalar conservation laws, Burgers equation.

**MSC2010:** 35F55, 35L65.

**Notations.** When  $1 \leq p \leq \infty$ , the natural norm in  $L^p(\mathbb{R}^n)$  is denoted  $\|\cdot\|_p$ , and the conjugate exponent of  $p$  is  $p'$ . The total space-time dimension is  $d = 1 + n$  and the coordinates are  $x = (t, y)$ . In the space of test functions,  $\mathcal{D}^+(\mathbb{R}^{1+n})$  is the cone of functions which take non-negative values. The partial derivative with respect to the coordinate  $y_j$  is  $\partial_j$ , while the time derivative is  $\partial_t$ . Various finite positive constants that depend only the dimension, but not upon the solutions

---

<sup>\*</sup>U.M.P.A., UMR CNRS-ENSL # 5669. École Normale Supérieure de Lyon. [denis.serre@ens-lyon.fr](mailto:denis.serre@ens-lyon.fr)

<sup>†</sup>Department of Mathematics, the University of Chicago. [luis@math.uchicago.edu](mailto:luis@math.uchicago.edu)

<sup>‡</sup>LS was partially supported by NSF grants DMS-1254332 and DMS-1764285.

of our PDE, are denoted  $c_d, c_{d,p}, c_{d,p,q}$ ; they usually differ from one inequality to another one. We denote  $C_0(0, +\infty)$  the space of continuous functions over  $(0, +\infty)$  which tend to zero at infinity. Mind that  $C(\mathbb{R}_+)$  is the space of bounded continuous functions over  $[0, +\infty)$ .

## 1 Introduction

Let us consider a scalar conservation law in  $1 + n$  dimensions

$$(1) \quad \partial_t u + \sum_{i=1}^n \partial_i f_i(u) = 0, \quad t > 0, y \in \mathbb{R}^n.$$

We complement this equation with an initial data

$$u(0, y) = u_0(y), \quad y \in \mathbb{R}^n.$$

The flux  $f(s) = (f_1(s), \dots, f_n(s))$  is a smooth vector-valued function of  $s \in \mathbb{R}$ . We recall the terminology that an entropy-entropy flux pair is a couple  $(\eta, q)$  where  $s \mapsto \eta(s)$  is a numerical function,  $s \mapsto q(s)$  a vector-valued function, such that  $q'(s) \equiv \eta'(s)f'(s)$ . The Kruzhkov's entropies and their fluxes form a one-parameter family:

$$\eta_a(s) = |s - a|, \quad q_a(s) = \operatorname{sgn}(s - a)(f(s) - f(a)).$$

Together with the affine functions, they span the cone of convex functions.

We recall that an *entropy solution* is a measurable function  $u \in L^1_{\text{loc}}([0, +\infty) \times \mathbb{R}^n)$  such that  $f(u) \in L^1_{\text{loc}}([0, +\infty) \times \mathbb{R}^n)$ , which satisfies the Cauchy problem in the distributional sense,

$$(2) \quad \int_0^\infty dt \int_{\mathbb{R}^n} (u \partial_t \phi + f(u) \cdot \nabla_y \phi) dy + \int_{\mathbb{R}^n} u_0(y) \phi(0, y) dy = 0, \quad \forall \phi \in \mathcal{D}(\mathbb{R}^{1+n}),$$

together with the entropy inequalities

$$(3) \quad \int_0^\infty dt \int_{\mathbb{R}^n} (\eta_a(u) \partial_t \phi + q_a(u) \cdot \nabla_y \phi) dy + \int_{\mathbb{R}^n} \eta_a(u_0(y)) \phi(0, y) dy \geq 0, \quad \forall \phi \in \mathcal{D}^+(\mathbb{R}^{1+n}), \forall a \in \mathbb{R}.$$

The theory of this Cauchy problem dates back to 1970, when S. Kruzhkov [10] proved that if  $u_0 \in L^\infty(\mathbb{R}^n)$ , then there exists one and only one entropy solution in the class

$$L^\infty(\mathbb{R}_+ \times \mathbb{R}^n) \cap C(\mathbb{R}_+; L^1_{\text{loc}}(\mathbb{R}^n)).$$

The parametrized family of operators  $S_t : u_0 \mapsto u(t, \cdot)$ , which map  $L^\infty(\mathbb{R}^n)$  into itself, form a semi-group. We warn the reader that  $S_t : L^\infty \rightarrow L^\infty$  is not continuous, because of the onset of shock waves. Likewise,  $t \mapsto u(t)$  is not continuous from  $\mathbb{R}_+$  into  $L^\infty(\mathbb{R}^n)$ .

This semi-group enjoys nevertheless nice properties. On the one hand, a comparison principle says that if  $u_0 \leq v_0$ , then  $S_t u_0 \leq S_t v_0$ . For instance, the solution  $u$  associated with the data  $u_0$  is majorized by the solution  $\bar{u}$  associated with the data  $(u_0)_+$ , the positive part of  $u_0$ . On another hand, if  $v_0 - u_0$  is integrable over  $\mathbb{R}^n$ , then  $S_t v_0 - S_t u_0$  is integrable too, and

$$(4) \quad \int_{\mathbb{R}^n} |S_t v_0 - S_t u_0|(y) dy \leq \int_{\mathbb{R}^n} |v_0 - u_0|(y) dy.$$

Finally,  $S_t$  maps  $L^p \cap L^\infty(\mathbb{R}^n)$  into itself, and the function  $t \mapsto \|S_t u_0\|_p$  is non-increasing.

Because of (4) and the density of  $L^1 \cap L^\infty(\mathbb{R}^n)$  in  $L^1(\mathbb{R}^n)$ , the family  $(S_t)_{t \geq 0}$  extends in a unique way as a continuous semi-group of contractions over  $L^1(\mathbb{R}^n)$ , still denoted  $(S_t)_{t \geq 0}$ . When  $u_0 \in L^1(\mathbb{R}^n)$  is unbounded, we are thus tempted to declare that  $u(t, y) := (S_t u_0)(y)$  is the *abstract solution* of the Cauchy problem for (1) with initial data  $u_0$ . At this stage, it is unclear whether  $(S_t)_{t \geq 0}$  can be defined as a semi-group over some  $L^p$ -space for  $p \in (1, \infty)$ , because the contraction property (4) occurs only in the  $L^1$ -distance, but in no other  $L^p$ -distance.

An alternate construction of  $(S_t)_{t \geq 0}$  over  $L^1(\mathbb{R}^n)$ , based upon the Generation Theorem for nonlinear semigroups, was done by M. Crandall [2], who pointed out that it is unclear whether  $u$  is an entropy solution, because the local integrability of the flux  $f(u)$  is not guaranteed<sup>1</sup>. The following question is therefore an important one:

*Identify the widest class of integrable initial data for which  $u$  is actually an entropy solution of (1).*

Our most complete results are about a special case, the so-called *multi-dimensional Burgers equation*

$$(5) \quad \partial_t u + \partial_j \frac{u^2}{2} + \cdots + \partial_n \frac{u^{n+1}}{n+1} = 0,$$

which is a paradigm of a genuinely non-linear conservation law. This equation was already considered by G. Crippa et al. [3], and more recently by L. Silvestre [17]. The particular flux in (5) is a prototype for genuinely nonlinear conservation laws, those which satisfy the assumption

$$(6) \quad \det(f'', \dots, f^{(n+1)}) \neq 0.$$

---

<sup>1</sup>Except of course in the case where  $f$  is globally Lipschitz.

The latter condition is a variant of the *non-degeneracy condition* at work in the kinetic formulation of the equation (1) ; see [12] or [13].

Our first result deals with dispersive estimates:

**Theorem 1.1** *Let  $1 \leq p \leq q \leq \infty$  be two exponents. Define two parameters  $\alpha, \beta(p, q)$  by*

$$(7) \quad \alpha(p, q) = \frac{h(q)}{h(p)}, \quad h(p) := 2 + \frac{dn}{p}$$

and

$$(8) \quad \beta(p, q) = h(q)(\delta(p) - \delta(q)), \quad \delta(p) := \frac{n}{2p + dn}.$$

*There exists a finite constant  $c_{d,p,q}$  such that for every initial data  $u_0 \in L^1 \cap L^\infty(\mathbb{R}^n)$ , the entropy solution  $u(t)$  of the scalar conservation law (5) satisfies*

$$(9) \quad \|u(t)\|_q \leq c_{d,p,q} t^{-\beta(p,q)} \|u_0\|_p^{\alpha(p,q)}, \quad \forall t > 0.$$

## Remarks

- The consistency of estimates (9) with the Hölder inequality is guaranted by the property that whenever  $\theta \in (0, 1)$ ,

$$(10) \quad \left( \frac{1}{q} = \frac{1-\theta}{p} + \frac{\theta}{r} \right) \implies \begin{cases} \alpha(p, q) = 1 - \theta + \theta\alpha(p, r), \\ \beta(p, q) = \theta\beta(p, r). \end{cases}$$

- The consistency under composition  $(p, q) \wedge (q, r) \mapsto (p, r)$  is ensured by the rules

$$(11) \quad \alpha(p, r) = \alpha(p, q)\alpha(q, r) \quad \text{and} \quad \beta(p, r) = \beta(q, r) + \beta(p, q)\alpha(q, r)$$

- In one space dimension, (9) gives back well-know results, such as Theorem<sup>2</sup> 11.5.2 in [6].

---

<sup>2</sup>Mind that this statement contains a typo, as the choice  $r = 1 - \frac{1}{p}$  in Theorem 11.5.1 yields the exponent  $-\frac{1}{p+1}$  instead of  $-\frac{p}{p+1}$ .

Theorem 1.1 has several important consequences. An obvious one is that the extension of  $(S_t)_{t \geq 0}$  as a semi-group over  $L^1(\mathbb{R}^n)$  satisfies the above estimates with  $p = 1$  :

**Corollary 1.1** *If  $u_0 \in L^1(\mathbb{R}^n)$  and  $t > 0$ , then  $S_t u_0 \in \bigcap_{1 \leq q \leq \infty} L^q(\mathbb{R}^n)$  and we have*

$$\|S_t u_0\|_q \leq c_{d,q} t^{-\kappa/q'} \|u_0\|_1^{1-\nu/q'}, \quad \forall q \in [1, \infty],$$

where the exponents are given in terms of

$$\kappa = 2 \frac{d-1}{d^2-d+2} \quad \text{and} \quad \nu = \frac{d(d-1)}{d^2-d+2}.$$

The next one is that the Cauchy problem is solvable for data taken in  $L^p(\mathbb{R}^n)$  for arbitrary exponent  $p \in [1, \infty]$ . In particular, it solves Crandall's concern.

**Theorem 1.2** *Let  $p \in [1, \infty)$  be given. For every  $t \geq 0$ , the operator  $S_t : L^1 \cap L^\infty(\mathbb{R}^n) \rightarrow L^1 \cap L^\infty(\mathbb{R}^n)$  admits a unique continuous extension  $S_t : L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ .*

*The family  $(S_t)_{t \geq 0}$  is a continuous semi-group over  $L^p(\mathbb{R}^n)$ . If  $u_0 \in L^p(\mathbb{R}^n)$ , the function  $u(t, y)$  defined by  $u(t) = S_t u_0$  is actually an entropy solution of the Cauchy problem for (5) with initial data  $u_0$ .*

*Finally,  $S_t(L^p(\mathbb{R}^n))$  is contained in  $\bigcap_{p \leq q \leq \infty} L^q(\mathbb{R}^n)$  and the estimates (9) are valid for every data  $u_0$  in  $L^p(\mathbb{R}^n)$ .*

The proof of Theorem 1.1 will be done in two steps. The first one consists in establishing the estimate (9) when  $q = p^*$  is given by the formula

$$p^* = d \left( 1 + \frac{p}{n} \right).$$

To this end, we apply Compensated Integrability to a suitable symmetric tensor, whose row-wise divergence is a bounded measure with controlled mass. This argument involves the theory recently developed by the first author in [14, 15]. The second step is an iteration in De Giorgi's style, based on the preliminary work [17] by the second author ; see also the original paper by E. De Giorgi [7] or the review paper by A. Vasseur [20]. This technique allows us to establish an  $L^\infty$ -estimate, which extends the dispersive estimate to  $q = +\infty$ . Then using the Hölder inequality, we may interpolate between this result and the decay of  $t \mapsto \|u(t)\|_p$ , and treat every exponent  $q > p$ .

We notice that the symmetric tensor mentioned above extends to a multi-dimensional context the one already used when  $n = 1$  by L. Tartar [19] to prove the compactness of the semi-group, and by F. Golse [8] (see also [9]) to prove some kind of regularity.

**Previous dispersive estimates.** In one space dimension  $n = 1$ , (5) reduces to the original Burgers equation. Its Kruzhkov solution satisfies the Oleinik inequality  $\partial_y u \leq \frac{1}{t}$ , which does not involve the initial data at all. Ph. B enilan & M. Crandall [1] proved

$$(12) \quad TV \left( \frac{u(t)^2}{2} \right) \leq \frac{2\|u_0\|_1}{t},$$

by exploiting the homogeneity of the flux  $f(s) = \frac{s^2}{2}$ . Inequality (12) implies an estimate

$$(13) \quad \|u(t)\|_\infty \leq 2\sqrt{\frac{2\|u_0\|_1}{t}},$$

which is a particular case of Corollary 1.1 in this simplest case.

C. Dafermos [5] proved a general form of (12) in situations where the flux  $f$  may have one inflexion point and the data  $u_0$  has bounded variations, by a clever use of the generalized backward characteristics. His argument involves the order structure of the real line. Backward characteristics are not unique in general. Given a base point  $(x^*, t^*)$  in the upper half-plane, one has to define and analyse the minimal and the maximal ones. The description of backward characteristics seems to be much more complicated in higher space dimensions, and Dafermos' strategy has not been applied successfully beyond the 1-D case.

**Enhanced decay.** Because of a scaling property which will be described in the next section, the dispersion (9) is optimal, as long as we involve only the  $L^p$ -norms, and we exclude any extra information about the initial data. It is however easy to obtain a better decay as time  $t$  goes to infinity. Let us give one example, by taking an initial data  $u_0$  such that

$$0 \leq u_0(y) \leq v_0(y_1), \quad v_0 \in L^1(\mathbb{R}).$$

By the maximum principle, we have  $u(t, y) \leq v(t, y_1)$ , where  $v$  is the solution of the 1-dimensional Burgers equation associated with the initial data  $v_0$ . We have therefore

$$\|u(t)\|_\infty \leq 2\sqrt{\frac{2\|v_0\|_1}{t}},$$

where the decay rate  $t^{-\frac{1}{2}}$  is independent of the space dimension. In particular this decay is faster than that given by Corollary 1.1 when  $n \geq 3$ .

The way this faster decay is compatible with the optimality of (9) is well explained by a study of the growth of the support of the solution. In the most favorable case where the data

$u_0$  is bounded with compact support, the argument above yields  $\|u(t)\|_\infty = O((1+t)^{-1/2})$ . It is easy to infer that the width of  $\text{Supp}(u(t))$  in the  $y_1$ -direction expands as  $O(\sqrt{t})$  (one might have used the comparison with the solution  $v$  above). Likewise, the width in the  $y_2$ -direction is an  $O(\log t)$  and that in the other  $y_k$ -directions remains bounded because

$$\int_0^\infty (1+t)^{-\frac{k}{2}} dt < \infty.$$

On the contrary, if  $u_0 \in L^1(\mathbb{R}^n)$  has compact support but is not bounded by an integrable fonction  $v_0(y_1)$  as above, Corollary 1.1 gives only  $\|u(t)\|_\infty = O(t^{-\kappa})$ . It turns out that  $n\kappa \geq 1$  when  $n \geq 2$ , and therefore

$$\int_0^\infty t^{-n\kappa} dt = +\infty.$$

This suggest that the width of the support in the  $y_n$ -direction is immediately infinite: the support of  $u(t)$  is unbounded for every  $t > 0$ . The solution has a tail in the last direction, and this tail is responsible for a slow  $L^\infty$ -decay, at rate  $t^{-\kappa}$  instead of  $t^{-\frac{1}{2}}$ .

This analysis suggests in particular that the *fundamental solution*  $U_m$ , if it exists, should have an unbounded support in the space variable when  $n \geq 2$ . The terminology denotes an entropy solution of (5), say a non-negative one, with the property that

$$U_m(t) \xrightarrow{t \rightarrow 0^+} m \delta_{y=0}$$

in the vague sense of bounded measures. In particular,

$$\int_{\mathbb{R}^n} U_m(t, y) dy \equiv m.$$

This behaviour is in strong contrast with the one-dimensional situation, where

$$U_m(t, y) = \frac{y}{t} \mathbf{1}_{(0, \sqrt{2mt})}$$

is compactly supported at every time.

The existence of a fundamental solution is left as an open problem. It should play an important role in the time-asymptotic analysis of entropy solutions of finite mass. This asymptotics has been known in one-space dimension since the seminal works by P. Lax [11] and C. Dafermos [4].

**Preliminary works.** The authors posted, separately, recent preprints on this subject in ArXiv database, see [16, 18]. The present paper supersedes both of them.

**Outline of the article.** We prove a special case of the dispersive estimate (9), that for the pairs  $(p, p^*)$ , in Section 2. We treat the case  $(p, \infty)$  in Section 3. This allows us to extend the (9) to every pair  $(p, q)$  with  $p \leq q$ . The construction of the semi-group over every  $L^p$ -space is done in Section 4. We show in Section 5 how these ideas adapt to a scalar equation when the fluxes  $f_j$  are monomials. The last section describes how the first argument, which involves Compensated Integrability, can be adapted to conservation laws with arbitrary flux.

**Acknowledgements.** We are indebted to C. Dafermos, who led us to collaborate.

## 2 Dispersive estimate ; the case $(p, p^*)$

To begin with, we recall that the Burgers equation enjoys an exceptional one-parameter transformation group, a fact already noted in [17] : Let  $u$  be an entropy solution of the Cauchy problem for (5) and  $\lambda$  be a positive constant. Then the function

$$v(t, y) = \frac{1}{\lambda} u(t, \lambda y_1, \dots, \lambda^n y_n)$$

is an entropy solution associated with the initial data

$$v_0(y) = \frac{1}{\lambda} u_0(\lambda y_1, \dots, \lambda^n y_n).$$

The following identities will be used below:

$$(14) \quad \int_0^\tau dt \int_{\mathbb{R}^n} v(t, y)^q dy = \lambda^{-q - \frac{d(d-1)}{2}} \int_0^\tau dt \int_{\mathbb{R}^n} u(t, y)^q dy,$$

$$(15) \quad \int_{\mathbb{R}^n} v_0(y)^q dy = \lambda^{-q - \frac{d(d-1)}{2}} \int_{\mathbb{R}^n} u_0(y)^q dy.$$

Let  $u_0^\pm$  be the positive and negative parts of the initial data:  $u_0^- \leq u_0 \leq u_0^+$  with  $u_0(x) \in \{u_0^-(x), u_0^+(x)\}$  everywhere. Denote  $u_\pm$  the entropy solutions associated with the data  $u_0^\pm$ . By the maximum principle, we have  $u_- \leq u \leq u_+$  everywhere. Because of  $\|u(t)\|_q \leq \|u_-(t)\|_q + \|u_+(t)\|_q$  and  $\|u_0\|_p = (\|u_0^-\|_p^p + \|u_0^+\|_p^p)^{1/p}$ , it suffices to prove the estimate for  $u_\pm$ , that is for initial data that are *signed*. And since  $v(t, y) = -u(t, -y_1, y_2, \dots, (-1)^n y_n)$  is the entropy solution associated with  $v_0(y) = -u_0(-y_1, y_2, \dots, (-1)^n y_n)$ , it suffices to treat the case of a non-negative initial data.

We therefore suppose from now on that  $u_0 \in L^1 \cap L^\infty(\mathbb{R}^n)$  and  $u_0 \geq 0$ , so that  $u \geq 0$  over  $\mathbb{R}_+ \times \mathbb{R}^n$ . We wish to estimate  $\|u(t)\|_q$  in terms of  $\|u_0\|_p$  when  $q = p^* = d(1 + \frac{p}{n})$ . We point out that  $p^* > p$ .

## 2.1 A Strichartz-like inequality

If  $a \in \mathbb{R}$ , we define a symmetric matrix

$$M(a) = \left( \frac{a^{i+j+p}}{i+j+p} \right)_{0 \leq i, j, \leq n}.$$

Remarking that

$$M(a) = \int_0^a V(s) \otimes V(s) s^{p-1} ds, \quad V(s) = \begin{pmatrix} 1 \\ \vdots \\ s^n \end{pmatrix},$$

we obtain that  $M(a)$  is positive definite whenever  $a > 0$ . Obviously,

$$\det M(a) = H_{d,p} a^{d(p+d-1)} = H_{d,p} a^{np^*},$$

where

$$H_{d,p} = \left\| \frac{1}{i+j+p} \right\|_{0 \leq i, j, \leq n} > 0$$

is a Hilbert-like determinant.

Let us form the symmetric tensor

$$T(t, y) = M(u(t, y)),$$

with positive semi-definite values. Its row of index  $i$  is formed of  $(\eta_{i+p}(u), q_{i+p}(u))$ , an entropy-flux pair where  $\eta_r(s) = \frac{|s|^r}{r}$  is convex. In the special case where  $p = 1$  and  $i = 0$ , it is divergence-free because of (5) itself. Otherwise, it is not divergence-free in general, although it is so wherever  $u$  is a classical solution. But the entropy inequality tells us that the opposite of its divergence is a non-negative, hence bounded measure,

$$\mu_r = -\operatorname{div}_{t,y}(\eta_r(u), q_r(u)) \geq 0.$$

The total mass of  $\mu_r$  over a slab  $(0, \tau) \times \mathbb{R}^n$  is given by

$$\|\mu_r\| = \int_{\mathbb{R}^n} \eta_r(u_0(y)) dy - \int_{\mathbb{R}^n} \eta_r(u(\tau, y)) dy \leq \int_{\mathbb{R}^n} \frac{u_0(y)^r}{r} dy.$$

Since the latter bound does not depend upon  $\tau$ ,  $\mu_r$  is actually a bounded measure over  $\mathbb{R}_+ \times \mathbb{R}^n$ .

We conclude that the row-wise divergence of  $T$  is a (vector-valued) bounded measure, whose total mass is bounded above by

$$\sum_{j=0}^n \int_{\mathbb{R}^n} \frac{u_0(y)^{j+p}}{j+p} dy.$$

We may therefore apply Compensated Integrability (Theorems 2.2 and 2.3 of [15]) to the tensor  $T$ , that is

$$\int_0^\tau dt \int_{\mathbb{R}^n} (\det T)^{\frac{1}{d-1}} dy \leq c_d \left( \|T_{0\bullet}(0, \cdot)\|_1 + \|T_{0\bullet}(\tau, \cdot)\|_1 + \|\operatorname{Div}_{t,y} T\|_{\mathcal{M}((0,\tau) \times \mathbb{R}^n)} \right)^{\frac{d}{d-1}}.$$

Because of

$$\|T_{0\bullet}(t, \cdot)\|_1 = \sum_{j=0}^n \int_{\mathbb{R}^n} \frac{u(t, y)^{j+p}}{j+p} dy \leq \sum_{j=0}^n \int_{\mathbb{R}^n} \frac{u_0(y)^{j+p}}{j+p} dy,$$

we deduce

$$(16) \quad \int_0^\tau dt \int_{\mathbb{R}^n} u^{p*} dy \leq c_{d,p} \left( \sum_{j=0}^n \int_{\mathbb{R}^n} u_0(y)^{j+p} dy \right)^{\frac{d}{d-1}}.$$

Again, the right-hand side does not depend upon  $\tau$ , thus the inequality above is true also for  $\tau = +\infty$ .

The only flaw in the estimate (16) is the lack of homogeneity of its right-hand side. To recover a well-balanced inequality, we use the scaling, in particular the formulæ (15). Applying (16) to the pair  $(v, v_0)$  instead, we get a parametrized inequality

$$\left( \int_0^\infty dt \int_{\mathbb{R}^n} u^{p*} dy \right)^{\frac{d-1}{d}} \leq c_d \lambda^{\frac{d-1}{2}} \sum_{j=0}^n \lambda^{-j} \int_{\mathbb{R}^n} u_0(y)^{j+p} dy,$$

where  $\lambda > 0$  is up to our choice. In order to minimize the right-hand side, we select the value

$$\lambda = \left( \int_{\mathbb{R}^n} u_0(y)^{n+p} dy / \int_{\mathbb{R}^n} u_0(y)^p dy \right)^{\frac{1}{n}}.$$

The extreme terms, for  $j = 0$  or  $n$ , contribute on an equal foot with

$$\left( \int_{\mathbb{R}^n} u_0(y)^{n+p} dy \right)^{\frac{1}{2}} \left( \int_{\mathbb{R}^n} u_0(y)^p dy \right)^{\frac{1}{2}}.$$

The other ones, which are

$$\left( \int_{\mathbb{R}^n} u_0(y)^{n+p} dy / \int_{\mathbb{R}^n} u_0(y)^p dy \right)^{\frac{1}{2} - \frac{j}{d-1}} \int_{\mathbb{R}^n} u_0^{j+p} dy,$$

are bounded by the same quantity, because of Hölder inequality. We end therefore with the fundamental estimate of Strichartz style

$$(17) \quad \left( \int_0^\infty \int_{\mathbb{R}^n} u^{p^*} dy dt \right)^{\frac{d-1}{d}} \leq c_d \left( \int_{\mathbb{R}^n} u_0(y)^{p+n} dy \right)^{\frac{1}{2}} \left( \int_{\mathbb{R}^n} u_0(y)^p dy \right)^{\frac{1}{2}}.$$

## 2.2 Proof of estimate (9)

We shall contemplate (17) as a differential inequality. To the end, we define

$$X(t) := \int_{\mathbb{R}^n} u^{p^*} dy = \|u(t)\|_{p^*}^{p^*}$$

Noticing that  $p+n$  is less than  $p^*$ , and using Hölder inequality, we get

$$\int_{\mathbb{R}^n} |w|^{p+n} dy \leq \left( \int_{\mathbb{R}^n} |w|^p dy \right)^a \left( \int_{\mathbb{R}^n} |w|^{p^*} dy \right)^b$$

for

$$a = \frac{p+n}{p+dn}, \quad b = \frac{n^2}{p+dn}.$$

The inequality (17) implies therefore

$$\left( \int_0^\infty X(t) dt \right)^{\frac{2n}{d}} \leq c_d \|u_0\|_p^{p(1+a)} X(0)^b.$$

Considering the solution  $w(t, y) = u(t + \tau, y)$ , whose initial data is  $u(\tau, \cdot)$ , we also have

$$(18) \quad \left( \int_\tau^\infty X(t) dt \right)^{\frac{2n}{db}} \leq c_d \|u(\tau)\|_p^{p \frac{1+a}{b}} X(\tau) \leq c_d \|u_0\|_p^{p \frac{1+a}{b}} X(\tau).$$

Let us denote

$$Y(\tau) := \int_\tau^\infty X(t) dt.$$

We recast (18) as

$$Y^\rho + c_d \|u_0\|_p^\mu Y' \leq 0, \quad \rho := \frac{2n}{db} \quad \mu := p \frac{1+a}{b}.$$

Remark that  $\rho = 2 \frac{p+dn}{dn} > 2$ . Multiplying by  $Y^{-\rho}$  and integrating, we infer

$$t + c_d \|u_0\|_p^\mu Y(0)^{1-\rho} \leq c_d \|u_0\|_p^\mu Y(t)^{1-\rho}.$$

This provides a first decay estimate

$$Y(t) \leq c_d \|u_0\|_p^{\frac{\mu}{\rho-1}} t^{-\frac{1}{\rho-1}}.$$

Remarking that  $t \mapsto X(t)$  is a non-increasing function, so that

$$\frac{\tau}{2} X(\tau) \leq Y\left(\frac{\tau}{2}\right),$$

we deduce the ultimate decay result

$$X(t) \leq c_d \|u_0\|_p^{\frac{\mu}{\rho-1}} t^{-\frac{\rho}{\rho-1}}.$$

Restated in terms of a Lebesgue norm of  $u(t)$ , it says

$$(19) \quad \|u(t)\|_{p^*} \leq c_d \|u_0\|_p^{\alpha(p,p^*)} t^{-\beta(p,p^*)},$$

where  $\alpha(p, q)$  and  $\beta(p, q)$  are given in (7) and (8). This is a special case of (9).

### 3 General pairs $(p, q)$ where $p < q \leq \infty$

Because of (10) and of the Hölder inequality, it will be enough to prove (9) when  $q = +\infty$ . Once again, it is sufficient to treat the case of non-negative data / solutions.

#### 3.1 An estimate for $(u - \ell)_+$

Let  $\ell > 0$  be a given number. We denote  $w_\ell$  the entropy solution of (5) associated with the initial data  $(u_0 - \ell)_+ + \ell = \max\{u_0, \ell\}$ . The function  $z_\ell := w_\ell - \ell$  is an entropy solution of a modified conservation law

$$\partial_t z_\ell + \sum_{k=1}^n \partial_k \frac{(z_\ell + \ell)^{k+1}}{k+1} = 0.$$

This is not exactly the Burgers equation for  $z_\ell$ . However the  $(n+2)$ -uplet  $(1, X + \ell, \dots, \dots, \frac{(X+\ell)^{n+1}}{n+1})$  is a basis of  $\mathbb{R}_{n+1}[X]$ . We pass from this basis to  $(1, X, \dots, \frac{X^{n+1}}{n+1})$  by a triangular matrix with unit diagonal. There exists therefore a change of coordinates

$$\begin{pmatrix} t \\ y' \end{pmatrix} = P \begin{pmatrix} t \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \vdots & Q \end{pmatrix} \begin{pmatrix} t \\ y \end{pmatrix},$$

where  $Q$  is a unitriangular matrix, such that  $z_\ell$  obeys the Burgers equation in the new coordinates:

$$\frac{\partial z_\ell}{\partial t} + \sum_{k=1}^n \frac{\partial}{\partial y'_k} \frac{z_\ell^{k+1}}{k+1} = 0.$$

We may therefore apply (19) to  $z_\ell$  :

$$\left( \int_{\mathbb{R}^n} z_\ell(t, y')^{p^*} dy' \right)^{\frac{1}{p^*}} \leq c_d \left( \int_{\mathbb{R}^n} z_\ell(0, y')^p dy' \right)^{\frac{\alpha(p, p^*)}{p}} t^{-\beta(p, p^*)}.$$

Remarking that the time variable is unchanged, and the Jacobian of the change of variable  $y \mapsto y'$  at fixed time equals one, we have actually

$$\|z_\ell(t)\|_{p^*} \leq c_d \|z_\ell(0)\|_p^{\alpha(p, p^*)} t^{-\beta(p, p^*)}.$$

Finally, the maximum principle tells us that  $u \leq w_\ell$ . The inequality above is therefore an estimate of the positive part of  $u - \ell$  :

$$(20) \quad \|(u - \ell)_+(t)\|_{p^*} \leq c_d \|(u_0 - \ell)_+\|_p^{\alpha(p, p^*)} t^{-\beta(p, p^*)}.$$

### 3.2 An iteration à la De Giorgi

We now prove the  $L^p$ - $L^\infty$  estimate, in the special case where  $\|u_0\|_p = 1$ . We recall that  $u_0$  is non-negative.

For the moment, we fix an arbitrary constant  $B > 0$ , which we will choose large enough in the end of the proof. Then we define the following sequences for  $k \in \mathbb{N}$  :

$$t_k = 1 - 2^{-k}, \quad \ell_k = B t_k, \quad w_k = (u - \ell_k)_+, \quad a_k = \|w_k(t_k)\|_p.$$

Remark that the sequences  $\ell_k$  and  $w_k$  are increasing and decreasing, respectively. Since  $t_0 = 0$ , we have  $a_0 = \|u_0\|_p = 1$ .

For each value of  $k$ , we apply (20) in order to estimate  $\|w_{k+1}(t_{k+1})\|_{p^*}$  in terms of  $\|w_{k+1}(t_k)\|_p$ . For the sake of simplicity, we write  $\alpha, \beta$  for  $\alpha(p, p^*)$  and  $\beta(p, p^*)$ . We get

$$\|w_{k+1}(t_{k+1})\|_{p^*} \leq c_{d,p} \|w_{k+1}(t_k)\|_p^\alpha (t_{k+1} - t_k)^{-\beta} = c_{d,p} 2^{\beta(k+1)} \|w_{k+1}(t_k)\|_p^\alpha \leq c_{d,p} 2^{\beta(k+1)} a_k^\alpha.$$

With Hölder inequality, we have also

$$a_{k+1} = \|w_{k+1}(t_{k+1})\|_p \leq \|w_{k+1}(t_{k+1})\|_{p^*} \|\mathbf{1}_{\{y: w_{k+1}(t_{k+1}, y) > 0\}}\|_r$$

where

$$\frac{1}{p} = \frac{1}{p^*} + \frac{1}{r}.$$

Remark that  $r > 1$ . Combining both inequalities, we obtain

$$a_{k+1} \leq c_{d,p} 2^{\beta(k+1)} a_k^\alpha |\{y : w_{k+1}(t_{k+1}, y) > 0\}|^{\frac{1}{r}}.$$

Observing that  $w_{k+1} > 0$  implies  $w_k > B2^{-k-1}$ , we infer

$$a_{k+1} \leq c_{d,p} 2^{\beta(k+1)} a_k^\alpha \left| \{y : w_k(t_{k+1}, y) > B2^{-k-1}\} \right|^{\frac{1}{r}}.$$

We now use Chebychev Inequality

$$\left| \{y : w_k(t_{k+1}, y) > B2^{-k-1}\} \right|^{\frac{1}{p}} \leq B^{-1} 2^{k+1} \|w_k(t_{k+1})\|_p \leq B^{-1} 2^{k+1} \|w_k(t_k)\|_p$$

to deduce

$$a_{k+1} \leq c_{d,p} B^{-\frac{p}{r}} 2^{(\beta + \frac{p}{r})(k+1)} a_k^{\alpha + \frac{p}{r}} = C2^{Ck} a_k^{1+\delta} B^{-\gamma}.$$

We have set  $\delta = \alpha - \frac{p}{p^*}$  and  $\gamma = \frac{p}{r}$ .

By a direct computation, we verify that  $\delta$  is positive:

$$\alpha - \frac{p}{p^*} = \frac{p^* h(p^*) - p h(p)}{p^* h(p)} = 2 \frac{p^* - p}{p^* h(p)} > 0.$$

The sequence  $b_k := B^{-\frac{\gamma}{\delta}} a_k$ , which starts with  $b_0 = B^{-\frac{\gamma}{\delta}}$ , satisfies therefore a recurrence relation

$$b_{k+1} \leq C2^{Ck} b_k^{1+\delta}.$$

It is known that if  $b_0$  is small enough, that is if  $B$  is large enough, then  $b_k \rightarrow 0+$  as  $k \rightarrow +\infty$ . Equivalently,  $a_k \rightarrow 0+$ .

We have therefore found a constant  $B > 0$  such that

$$\|(u - \ell_k)_+(1)\|_p \leq \|(u - \ell_k)_+(t_k)\|_p = a_k \rightarrow 0+.$$

Since  $\ell_k \rightarrow B$ , this means exactly that  $\|u(1)\|_\infty \leq B$ .

### 3.3 End of the proof of dispersive estimates

Let  $u_0 \in L^1 \cap L^\infty(\mathbb{R}^n)$  be non-negative. For two positive parameters  $\lambda, \mu$ , the entropy solution associated with the data

$$v_0(y) = \frac{1}{\lambda} u_0(\mu \lambda y_1, \dots, \mu \lambda^n y_n)$$

is the function

$$v(t, y) = \frac{1}{\lambda} u(\mu t, \mu \lambda y_1, \dots, \mu \lambda^n y_n).$$

If

$$(21) \quad \lambda^{p + \frac{n(n+1)}{2}} \mu^n = \int_{\mathbb{R}^n} u_0(y)^p dy,$$

then  $\|v_0\|_p = 1$  and we may apply the previous paragraph:  $\|v(1)\|_\infty \leq B$ . In terms of  $u$ , this writes

$$\|u(\mu)\|_\infty \leq B\lambda.$$

Eliminating  $\lambda$  with (21), this gives

$$\|u(\mu)\|_\infty \leq B \left( \mu^{-n} \|u_0\|_p^p \right)^{\frac{2}{n^2 + n + 2p}},$$

which is nothing but the dispersive estimate (9) for  $q = +\infty$ .

There remains to pass from  $q = +\infty$  to every  $q \in [p, +\infty]$ . We do that by applying the Hölder inequality. Writing

$$\frac{1}{q} = \frac{1-\theta}{p} + \frac{\theta}{\infty},$$

we have

$$\|u(t)\|_q \leq \|u(t)\|_p^{1-\theta} \|u(t)\|_\infty^\theta \leq \|u_0\|_p^{1-\theta} \left( B t^{-\beta(p, \infty)} \|u_0\|_p^{\alpha(p, \infty)} \right)^\theta.$$

We conclude by using the relations (10).

## 4 The $L^p$ -semi-group for finite exponents

We now prove Theorem 1.2. We start with a remark about  $L^p$ -spaces.

**Lemma 4.1** *Let  $a \in L^p(\mathbb{R}^n)$  be given. There exists a sequence  $(b_m)_{m \geq 0}$  in  $(L^p \cap L^\infty)(\mathbb{R}^n)$ , converging towards  $a$  in  $L^p(\mathbb{R}^n)$ , such that  $b_m - a \in L^1(\mathbb{R}^n)$  and*

$$\lim_{m \rightarrow +\infty} \|b_m - a\|_1 = 0.$$

*Proof*

Recall that

$$L^p(\mathbb{R}^n) = (L^1 \cap L^p)(\mathbb{R}^n) + (L^p \cap L^\infty)(\mathbb{R}^n).$$

Decomposing our function as  $a = a_1 + a_\infty$  where

$$a_1 \in (L^1 \cap L^p)(\mathbb{R}^n), \quad a_\infty \in (L^p \cap L^\infty)(\mathbb{R}^n),$$

we may form the sequence of bounded functions  $b_m := a_\infty + \pi_m \circ a_1$ , where  $\pi_m$  is the projection from  $\mathbb{R}$  onto the interval  $[-m, m]$ . Because of

$$\|b_m\|_p \leq \|a_\infty\|_p + \|\pi_m \circ a_1\|_p \leq \|a_\infty\|_p + \|a_1\|_p,$$

this sequence is bounded in  $L^p(\mathbb{R}^n)$ . In addition  $b_m - a = \pi_m \circ a_1 - a_1 \in L^1 \cap L^p(\mathbb{R}^n)$ , and

$$\|b_m - a\|_1 = \|\pi_m \circ a_1 - a_1\|_1 \xrightarrow{m \rightarrow +\infty} 0, \quad \|b_m - a\|_p = \|\pi_m \circ a_1 - a_1\|_p \xrightarrow{m \rightarrow +\infty} 0.$$

■

Let  $u_0 \in L^p(\mathbb{R}^n)$  be given. In order to define  $S_t u_0$ , we consider a sequence  $b_m$  that approximates  $u_0$  in the sense of Lemma 4.1. Remark that we do not care about the construction of  $b_m$ , as we only use the properties stated in the Lemma.

To begin with,  $u_m(t) := S_t b_m$  is well-defined and belongs to  $L^\infty(\mathbb{R}^n)$ . Because of (9), we have

$$(22) \quad \|u_m(t)\|_q \leq c_{d,p,q} \|b_m\|_p^{\alpha(p,q)} t^{-\beta(p,q)} \leq C_{p,q}(u_0) t^{-\beta(p,q)}.$$

The sequence  $(u_m)_{m>0}$  is thus bounded in  $C_0(\tau, \infty; L^q(\mathbb{R}^n))$  for every  $q \in [p, \infty)$  and every  $\tau > 0$ .

The contraction property gives us

$$\|u_m(t) - u_\ell(t)\|_1 \leq \|b_m - b_\ell\|_1 \xrightarrow{m, \ell \rightarrow +\infty} 0.$$

Let  $r, q$  be exponents satisfying  $p \leq r < q < \infty$ . By Hölder inequality, we have

$$\|u_m(t) - u_\ell(t)\|_r \leq \|u_m(t) - u_\ell(t)\|_1^\theta (\|u_m(t)\|_q + \|u_\ell(t)\|_q)^{1-\theta},$$

where  $\theta \in (0, 1]$ . With (22), we infer that

$$\|u_m(t) - u_\ell(t)\|_r \xrightarrow{m, \ell \rightarrow +\infty} 0,$$

uniformly over  $(\tau, \infty)$ .

We have thus proved that  $(u_m)_{m>0}$  is a Cauchy sequence in  $C_0(\tau, \infty; L^r(\mathbb{R}^n))$ , hence is convergent in this space. If  $b'_m$  is another approximating sequence for  $u_0$ , and  $u'_m$  the corresponding solution of the Cauchy problem, we may form an approximating sequence  $c_m$  in the sense of Lemma 4.1, by alternating  $b_1, b'_1, b_2, b'_2, \dots$ . The sequence  $u_1, u'_1, u_2, u'_2, \dots$  will be convergent in the sense above. This shows that the limit of  $u_m$  does not depend upon the precise sequence  $(b_m)_{m>0}$  chosen above. Thus we may set

$$S_t u_0 := \lim_{m \rightarrow +\infty} u_m(t),$$

which defines a

$$u \in C_b(\mathbb{R}_+; L^p(\mathbb{R}^n)) \bigcap \bigcap_{p < r < \infty} C_0(0, +\infty; L^r(\mathbb{R}^n)).$$

There remains to prove that  $u$  is an entropy solution of (5). For this, we use the fact that  $u_m$  is itself an entropy solution, and the convergence stated above ensures that every monomial  $(u_m)^j$  in the flux  $f(u_m)$ , converges towards  $u^j$  in  $L^1_{\text{loc}}$ .

The fact that  $u(0) = u_0$  follows from  $u_m(0) = b_m$ , the  $L^p$ -convergence  $b_m \rightarrow u_0$ , and the uniform convergence  $u_m(t) \rightarrow u(t)$  in  $L^p(\mathbb{R}^n)$ .

## 5 Other “monomial” scalar conservation laws

We consider in this section conservation laws whose fluxes are monomial. Denoting  $m_k(s) = \frac{s^{k+1}}{k+1}$ , they bear the form

$$(23) \quad \partial_t u + \partial_1 m_{k_1}(u) + \dots + \partial_n m_{k_n}(u) = 0,$$

where  $0 < k_1 < \dots < k_n$  are integers. The time derivative may be written as well  $\partial_t m_{k_0}(u)$  with  $k_0 = 0$ .

As before, we may restrict to non-negative initial data  $u_0$  that belong to  $L^1 \cap L^\infty(\mathbb{R}^n)$ . Given an exponent  $p \geq 1$ , our symmetric tensor is  $T(t, y) = M(u(t, y))$  where now

$$M(a) := (m_{p+k_i+k_j-1}(a))_{0 \leq i, j \leq n}.$$

Notice that  $M(a)$  is symmetric, and its upper-left entry is  $\frac{a^p}{p}$ . Because of

$$M(a) = \int_0^a s^{p-1} V(s) \otimes V(s) ds, \quad V(s) := \begin{pmatrix} s^{k_0} \\ \vdots \\ s^{k_n} \end{pmatrix},$$

it positive definite whenever  $a > 0$ . We have

$$\det M(a) = \Delta(p, \vec{k}) a^N, \quad N = dp + 2K, \quad K := \sum_0^n k_i.$$

As above, the lines of  $T$  are made of entropy-entropy flux pairs of the equation (23). Its row-wise divergence is therefore a vector-valued bounded measure. Compensated integrability yields again an inequality

$$\left( \int_0^\infty dt \int_{\mathbb{R}^n} u(t, y)^Q dy \right)^{\frac{n}{d}} \leq c_{d,p,\vec{k}} \sum_{j=0}^n \int_{\mathbb{R}^n} u_0(y)^{p+k_j} dy, \quad Q := \frac{N}{n}.$$

The conservation law is invariant under the scaling

$$u \longmapsto v(t, y) := \frac{1}{\lambda} u(t, \lambda^{k_1} y_1, \dots, \lambda^{k_n} y_n).$$

Applying the estimate above to  $v$ , we obtain a parametrized inequality :

$$\left( \int_0^\infty dt \int_{\mathbb{R}^n} u(t, y)^Q dy \right)^{\frac{n}{d}} \leq c_{d,p,\vec{k}} \lambda^{\frac{K}{d}} \sum_{j=0}^n \lambda^{-k_j} \int_{\mathbb{R}^n} u_0(y)^{p+k_j} dy.$$

We now choose

$$\lambda = \left( \int_{\mathbb{R}^n} u_0(y)^{p+k_n} dy / \int_{\mathbb{R}^n} u_0(y)^p dy \right)^{\frac{1}{k_n}}$$

and obtain a Strichartz-like estimate:

$$\left( \int_0^\infty dt \int_{\mathbb{R}^n} u(t, y)^Q dy \right)^{\frac{n}{d}} \leq c_{d,p,\vec{k}} \left( \int_{\mathbb{R}^n} u_0(y)^{p+k_n} dy \right)^\theta \left( \int_{\mathbb{R}^n} u_0(y)^p dy \right)^{1-\theta}$$

where

$$\theta := \frac{K}{dk_n} \in (0, 1).$$

Applying this calculation to the interval  $(\tau, +\infty)$ , and using the decay of the  $L^p$ -norm, we infer

$$(24) \quad \left( \int_\tau^\infty dt \int_{\mathbb{R}^n} u(t, y)^Q dy \right)^{\frac{n}{d}} \leq c_{d,p,\vec{k}} \left( \int_{\mathbb{R}^n} u(\tau, y)^{p+k_n} dy \right)^\theta \left( \int_{\mathbb{R}^n} u_0(y)^p dy \right)^{1-\theta}$$

We may now continue the analysis with a Gronwall argument, provided  $p + k_n \in (p, Q]$ . We leave the interested reader to check the details. Our first dispersion estimate is

$$(25) \quad \|u(t)\|_Q \leq c_{d,p} t^{-\beta(p)} \|u_0\|_p^{\alpha(p)},$$

whenever  $p \geq nk_n - 2K$  (remark that for the Burgers equation, this restriction is harmless).

At this stage, it seems that we miss an argument in order to carry out the De Giorgi technique, because the conservation law satisfied by  $u - \ell$  will be a different one. Whether it can be done here and for general conservation laws is left for a future work. What we can do at least is to combine the estimates (25) in order to cover pairs  $(p, q)$  of finite exponents. For instance, starting from a pair  $(p, Q)$  as above and choosing  $p_1 = Q$ , we have a corresponding  $Q_1$  such that (25) applies with  $(p_1, Q_1)$  instead of  $(p, Q)$ . We infer

$$\|u(t)\|_{Q_1} \leq c_{d,Q} (t/2)^{-\beta(Q)} \|u(t/2)\|_Q^{\alpha(Q)} \leq c_{d,p} t^{-\beta(Q) - \alpha(Q)\beta(p)} \|u_0\|_p^{\alpha(p)\alpha(Q)}.$$

Because the iteration  $p \rightarrow Q$  defines a sequence which tends to  $+\infty$ , and using the Hölder inequality to fill the gaps, we deduce the dispersion inequalities for the monomial conservation law:

**Theorem 5.1** *For the scalar conservation law (23) with monomial fluxes, there exist finite constants  $c_{d,p,q}$  such that whenever  $p \geq nk_n - 2K$ ,  $q \in [p, \infty)$  and  $u_0 \in L^p \cap L^\infty(\mathbb{R}^n)$ , we have*

$$\|u(t)\|_q \leq c_{d,p,q} t^{-\beta(p,q)} \|u_0\|_p^{\alpha(p,q)}.$$

*The exponents are given by the formula*

$$\alpha(p, q) = \frac{h(q)}{h(p)}, \quad h(p) := 1 + \frac{K}{p} \quad \text{and} \quad \beta(p, q) = n \left( \frac{\alpha(p, q)}{p} - \frac{1}{q} \right).$$

As in the case of the Burgers equation, we can use these estimates in order to define the semi-group over  $L^p$ -spaces:

**Corollary 5.1** *The semi-group  $(S_t)_{t \geq 0}$  for equation (23) extends by continuity as a continuous semi-group over  $L^p(\mathbb{R}^n)$  for every  $p \in [1, +\infty)$  such that  $p \geq nk_n - 2K$ . It maps  $L^p(\mathbb{R}^n)$  into  $L^q(\mathbb{R}^n)$  for every  $q \in [p, \infty)$ . If  $u_0 \in L^p(\mathbb{R}^n)$ , then the function  $u(t, y) := (S_t u_0)(y)$  is an entropy solution with initial data  $u_0$ .*

## 6 Compensated integrability for general fluxes $f$

We consider now a multi-dimensional conservation law of the most general form (1). Following the ideas developped in the Burgers and monomial cases, we begin by considering a signed, bounded initial data:  $u_0 \in L^1 \cap L^\infty(\mathbb{R}^n)$ ,  $u_0 \geq 0$ . If  $a \in \mathbb{R}_+$ , we define a symmetric matrix

$$M_g(a) = \int_0^a g(s) Z'(s) \otimes Z'(s) ds,$$

where  $Z(s) = (f_0(s) = s, f_1(s), \dots, f_n(s))$  and  $g$  is some positive function. This matrix is positive definite under the non-degeneracy condition that  $Z([0, a])$  is not contained in an affine hyperplane. We denote

$$\Delta_g(a) := (\det M_g(a))^{\frac{1}{n}} \geq 0.$$

Let us define  $T(t, y) := M_\Phi(u(t, y))$ . Because of  $u \in L^\infty(\mathbb{R}_+; L^1 \cap L^\infty(\mathbb{R}^n))$ , the tensor  $T$  is integrable over  $(0, \tau) \times \mathbb{R}^n$ . Each row of  $T$  is made of entropy-entropy flux pairs  $(F_i, Q_i)$ . Since  $F_i$  might not be convex, we cannot estimate the measure  $\mu_i = -\partial_t F_i(u) - \operatorname{div}_y Q_i(u)$  directly by the integral of  $F_i(u_0)$ . To overcome this difficulty, we define a convex function  $\phi_g$  over  $\mathbb{R}_+$  by

$$\phi_g(0) = \phi'_g(0) = 0, \quad \phi''_g(s) = |F''(s)|,$$

where  $F = (F_0, \dots, F_n)$ . Remark that  $|F'| \leq \phi'_g$  and  $|F| \leq \phi_g$ . Let  $\Phi_g$  be the entropy flux associated with the entropy  $\phi_g$ . Then the measure  $\mathbf{v}_g := -\partial_t \phi_g(u) - \operatorname{div}_y \Phi_g(u)$  is non-negative and a bound of its total mass is as usual

$$\|\mathbf{v}_g\| \leq \int_{\mathbb{R}^n} \phi_g(u_0(y)) dy.$$

We now use the kinetic formulation of (1), a notion for which we refer to [13], Theorem 3.2.1. Recall the definition of the kinetic function  $\chi(\xi; a)$ , whose value is  $\operatorname{sgn} a$  if  $\xi$  lies between 0 and  $a$ , and is 0 otherwise. There exists a non-negative bounded measure  $m(t, y, \xi)$  such that the function  $w(t, y, \xi) = \chi(\xi; u(t, y))$  satisfies

$$\partial_t w + f'(\xi) \cdot \nabla_y w = \frac{\partial}{\partial \xi} m, \quad w(0, y; \xi) = \chi(\xi; u_0(y)).$$

If  $(\eta, q)$  is an entropy-entropy flux pair, then the measure  $\mu = -\partial_t \eta - \operatorname{div}_y q$  is given by

$$\mu = \int_{\mathbb{R}} \eta''(\xi) dm(\xi).$$

We deduce that the vector-valued measure  $\mu = (\mu_0, \dots, \mu_n)$  satisfies  $|\mu| \leq v_g$ . This yields the estimate

$$\|\mu\| \leq \int_{\mathbb{R}^n} \phi_g(u_0(y)) dy.$$

We may therefore apply the compensated integrability, which gives here

$$\int_0^\tau dt \int_{\mathbb{R}^n} \Delta_g(u(t, y)) dy \leq c_d \left( \|F(u_0)\|_1 + \|F(u(\tau))\|_1 + \int_{\mathbb{R}^n} \phi_g(u_0(y)) dy \right)^{1+\frac{1}{n}}.$$

Because of  $|F| \leq \phi_g$  and  $\|\phi_g(u(\tau))\|_1 \leq \|\phi_g(u_0)\|_1$ , we end up with an analog of (17)

$$(26) \quad \int_0^\infty dt \int_{\mathbb{R}^n} \Delta_g(u(t, y)) dy \leq c_d \|\phi_g(u_0)\|_1^{1+\frac{1}{n}}.$$

Whether (26) can be used to prove dispersive estimates depends of the amount of nonlinearity of the equation (1). We leave this question for a future work.

## References

- [1] Ph. B enilan, M. G. Crandall. Regularizing effects of homogeneous evolution equations. *Contributions to analysis and geometry (Baltimore, Md., 1980)*. Johns Hopkins Univ. Press, Baltimore, Md. (1981), pp 23–39.
- [2] M. Crandall. The semigroup approach to first order quasilinear equations in several space variables. *Israel J. Math.*, **12** (1972), pp 108–132.
- [3] G. Crippa, F. Otto, M. Westdickenberg. Regularizing effect of nonlinearity in multidimensional scalar conservation laws. *Transport equations and multi-D hyperbolic conservation laws*, Lect. Notes Unione Mat. Ital., **5**, Springer, Berlin, (2008), pp 77–128.
- [4] C. Dafermos. Characteristics in hyperbolic conservation laws. *Nonlinear Analysis and Mechanics: Heriot-Watt Symposium (Edinburgh 1976)*, Vol. I, pp 1–58, ed. R. J. Knops. Research Notes in Math., No 17, Pitman, London (1977).
- [5] C. Dafermos. Regularity and large time behaviour of solutions of a conservation law without convexity. *Proc. Royal Soc. Edinburgh*, **99A** (1985), pp 201–239.
- [6] C. Dafermos. *Hyperbolic conservation laws in continuum physics*. Grundlehren der mathematischen Wissenschaften vol. 325, 3rd ed. Springer-Verlag, Heidelberg (2010).

- [7] E. De Giorgi. Sulla differenziabilità e l'analiticità delle estremali degli integrali multipli regolari. *Mem. Accad. Sci. Torino. Cl. Sci. Fis. Mat. Nat. (3)*, **3** (1957), pp 25–43.
- [8] F. Golse. Nonlinear regularizing effect for conservation laws. In *Hyperbolic problems: theory, numerics and applications*. Proc. Sympos. Appl. Math. 67, Part 1, Amer. Math. Soc., Providence, RI, (2009), pp 73–92.
- [9] F. Golse, B. Perthame. Optimal regularizing effect for scalar conservation laws. *Rev. Mat. Iberoam.*, **29** (2013), pp 1477–1504.
- [10] S. Kružkov. First order quasilinear equations with several independent variables (in Russian). *Mat. Sbornik (N.S.)*, **81 (123)** (1970), pp 228–255.
- [11] P. Lax. Hyperbolic systems of conservation laws. *Comm. Pure Appl. Math.* **10** (1957), pp 537–566.
- [12] P.-L. Lions, B. Perthame, E. Tadmor. A kinetic formulation of multidimensional scalar conservation laws and related equations. *J. Amer. Math. Soc.*, **7** (1994), pp 169–191.
- [13] B. Perthame. *Kinetic formulation of conservation laws*, Oxford lecture series in Math. & its Appl. **21**. Oxford (2002).
- [14] D. Serre. Divergence-free positive symmetric tensors and fluid dynamics. *Annales de l'Institut Henri Poincaré (analyse non linéaire)*. **35** (2018), pp 1209–1234. <https://doi.org/10.1016/j.anihpc.2017.11.002>.
- [15] D. Serre. Compensated integrability. Applications to the Vlasov–Poisson equation and other models in mathematical physics. *Journal de Mathématiques Pures et Appliquées*. To appear.
- [16] D. Serre. Multi-dimensional scalar conservation laws with unbounded integrable initial data. Preprint arXiv:1807.10474.
- [17] L. Silvestre. Oscillation properties of scalar conservation laws. Preprint arXiv:1708.03401v3.
- [18] L. Silvestre. A dispersive estimate for the multidimensional Burgers equation. Preprint arXiv:1808.01220.
- [19] L. Tartar. Compensated compactness and applications to partial differential equations. *Nonlinear analysis and mechanics: Heriot-Watt Symposium*, Vol. IV, Res. Notes in Math., 39, Pitman (1979), pp 136–212.

- [20] A. Vasseur. The De Giorgi method for elliptic and parabolic equations and some applications. *Lectures on the analysis of nonlinear partial differential equations. Part 4*, 195–222, Morningside Lect. Math., 4, *Int. Press, Somerville, MA*, 2016.