



**HAL**  
open science

# Mixed determinants, Compensated Integrability and new *a priori* estimates in Gas dynamics

Denis Serre

► **To cite this version:**

Denis Serre. Mixed determinants, Compensated Integrability and new *a priori* estimates in Gas dynamics. 2022. ensl-03945122

**HAL Id: ensl-03945122**

**<https://hal-ens-lyon.archives-ouvertes.fr/ensl-03945122>**

Preprint submitted on 10 Feb 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Mixed determinants, Compensated Integrability and new *a priori* estimates in Gas dynamics

Denis Serre  
École Normale Supérieure de Lyon \*

*To Constantine Dafermos, with admiration, friendship and gratitude.*

## Résumé

We extend the scope of our recent Compensated Integrability theory, by exploiting the multi-linearity of the determinant map over  $\mathbf{Sym}_n(\mathbb{R})$ . This allows us to establish new *a priori* estimates for inviscid gases flowing in the whole space  $\mathbb{R}^d$ . Notably, we estimate the defect measure (Boltzman equation) or weighted spacial correlations of the velocity field (Euler system). As usual, our bounds involve only the total mass and energy of the flow.

**Keywords:** Div-BV tensors, mixed determinant, Schur complement, gas dynamics.

**2020 Mathematics subject classification:** 35Q31, 46N60, 46T99, 76N15

**Notations:** The unit sphere of  $\mathbb{R}^n$  is  $S_{n-1}$ , whose area is  $|S_{n-1}|$ . For  $1 \leq j \leq n$  and  $x \in \mathbb{R}^n$ ,  $\hat{x}_j \in \mathbb{R}^{n-1}$  denotes the vector in which the  $j$ th coordinate is omitted.

The exponent  $T$  indicates transposition. Given a vector  $Z$ , we set  $Z \otimes Z := ZZ^T$  the symmetric, rank-one matrix of entries  $z_i z_j$ . The cofactor matrix of an  $n \times n$  matrix  $M$  is  $\widehat{M}$ . We recall the identities  $M^T \widehat{M} = \widehat{M} M^T = (\det M) I_n$  and  $\det \widehat{M} = (\det M)^{n-1}$ . The space of  $n \times n$  symmetric matrices with real entries being  $\mathbf{Sym}_n$ , the cone of positive semi-definite ones is  $\mathbf{Sym}_n^+$ . Its interior is  $\mathbf{SPD}_n$ . The inequality  $A \succ B$  between symmetric matrices, means that  $A - B \in \mathbf{Sym}_n^+$ .

An inequality  $F(X) \leq_n G(X)$  means that there exists a constant  $C(n)$ , depending only upon the ambient dimension  $n$ , such that every  $X$  under consideration satisfies  $F(X) \leq C(n) \cdot G(X)$ .

---

\*. U.M.P.A., UMR CNRS-ENSL # 5669. 46 allée d'Italie, 69364 Lyon cedex 07. France.  
denis.serre@ens-lyon.fr

The total mass of a finite Radon measure  $\mu$  is  $\|\mu\|_{\mathcal{M}}$ . If  $\vec{\mu}$  is a vector of Radon measures, its Euclidian norm  $|\vec{\mu}|$  is still a Radon measure; we again write  $\|\vec{\mu}\|_{\mathcal{M}}$  for the mass of  $|\vec{\mu}|$ .

## 1 Introduction

We recall that a *symmetric tensor* over  $\mathbb{R}^n$  is a symmetric  $n \times n$  matrix  $A$ , whose entries are distributions,  $a_{ij} \in \mathcal{D}(\mathbb{R}^n)$ . The Divergence (with a capital letter in this context) operator associates with  $A$  a vector of distributions,

$$(\text{Div}A)_i = \sum_{j=1}^n \partial_j a_{ij}, \quad \forall i \in \llbracket 1, n \rrbracket.$$

Recall that if  $A$  is positive semi-definite, then the  $a_{ij}$ 's are Radon measures. Since they are absolutely continuous with respect to  $\text{Tr} A$ , and because  $\det^{\frac{1}{n}} : \mathbf{Sym}_n^+ \rightarrow \mathbb{R}$  is positively homogeneous of order 1, we can define unambiguously the Radon measure  $(\det A)^{\frac{1}{n}}$ .

**Definition 1.1** *The tensor  $A$  is said Div-BV over  $\mathbb{R}^n$  if its entries  $a_{ij}$ , as well as the coordinates  $(\text{Div}A)_i$ , are Radon measures with finite total masses.*

The fundamental statement of Compensated Integrability, established in [6, 7] reads as follows.

**Theorem 1.1 ([6, 7])** *Let  $A$  be a positive semi-definite Div-BV tensor over  $\mathbb{R}^n$ . Then  $(\det A)^{\frac{1}{n}} \in L^{\frac{n}{n-1}}(\mathbb{R}^n)$  and*

$$\int_{\mathbb{R}^n} (\det A)^{\frac{1}{n-1}} dx \leq c_n \|\text{Div}A\|_{\mathcal{M}}^{\frac{n}{n-1}}.$$

This inequality is sharp, reducing to the isoperimetric inequality when  $A = \chi_{\Omega} I_n$ .

Theorem 1.1 implies several variants that have been described in some other papers of ours. Of particular interest is a version taylorred for Cauchy problems in the whole space  $\mathbb{R}^d$ . There, a tensor is given in the strip  $Q_T = (0, T) \times \mathbb{R}^d$ , which is often Div-free. We set  $n = 1 + d$  and  $x = (t, y)$  where  $y$  is the space variable. The tensor is extended by  $0_n$  away from  $Q_T$ . This extension is Div-BV provided that the traces of the first column (that is the normal traces) at  $t = 0^+$  and  $t = T^-$  are finite measures.

For the most important example of inviscid gas dynamics, the tensor is given in  $Q_T$  by

$$\begin{pmatrix} \rho & \rho u^T \\ \rho u & \rho u \otimes u + pI_d \end{pmatrix}$$

where  $\rho \geq 0$  is the mass density,  $p \geq 0$  the pressure and  $u$  is the fluid velocity. The specific internal energy  $e \geq 0$  is given in terms of  $\rho, p$  through an equation of state, typically  $p = (\gamma - 1)\rho e$  for a perfect gas,  $\gamma > 1$  being the adiabatic constant. The trace condition is ensured by the assumption of finite total mass and total mechanical energy. The application of C.I. has thus led to the following estimate of the internal variables:

**Theorem 1.2 ([6, 7])** *Consider an admissible inviscid gas flow in  $Q_T$  of finite total mass and energy,*

$$M := \int_{\mathbb{R}^d} \rho(0, y) dy < +\infty, \quad E_0 := \int_{\mathbb{R}^d} \left( \frac{1}{2} \rho |u|^2 + \rho e \right) (0, y) dy < +\infty.$$

Then

$$(1) \quad \int_{Q_T} \rho^{\frac{1}{d}} p(t, y) dy dt \leq_d M^{\frac{1}{d}} \sqrt{ME_0}.$$

We recall that since the right-hand side is independent from  $T$ , (1) is valid when  $T = +\infty$ . The same remark holds true for all the statements below, unless otherwise specified.

The admissibility assumed in Theorem 1.2 is the fact the total mass is conserved and the total energy at positive times is bounded by that at time  $t = 0$ :

$$(2) \quad \forall t \in (0, T) \quad \int_{\mathbb{R}^d} \rho(t, y) dy = M, \quad \int_{\mathbb{R}^d} \left( \frac{1}{2} \rho |u|^2 + \rho e \right) (t, y) dy \leq E_0.$$

The strength of Estimate (1), when compared with (2.b), is the presence of an extra factor  $\rho^{\frac{1}{d}}$  in the integrand. The price to pay is the replacement of a supremum over  $t \in (0, T)$  by a time integral. Somehow, (1) plays the role of a Strichartz inequality for Gas dynamics.

The flaw of Theorem 1.2 is of course the lack of a corresponding estimate for the velocity field. This is unsatisfactory since the energy estimate (2.b) suggests that  $\rho |u|^2$  and  $\rho e$  (proportional to  $p$ ) are on the par, having the same physical dimension. We should expect a space-time estimate of  $\rho^{1+\frac{1}{d}} |u|^2$ , as a counterpart of (1).

The purpose of this paper is thus to establish complementary estimates for Gas dynamics. They have several features in common with (1). First, they concern space-time integrals. Second, the integrands contain an extra factor not present in the energy density. Third, the bounds are still given in terms of  $M$  and  $E_0$ . At last these estimates are invariant under Galilean transformations. Somehow, all these estimate might be viewed as *fundamental*, or universal; in particular, they do not depend upon the particular equation of state.

Most importantly, we fill the gap mentioned above, by handling the velocity field. Mind however that since we are only able to treat Galilean invariant quantities, the velocity enters in the integrals through correlations

$$\text{Cor}(u_0, \dots, u_d) = \det \begin{pmatrix} 1 & \cdots & 1 \\ u_0 & \cdots & u_d \end{pmatrix},$$

where  $u_j(t, y) := u(t, y + h_j)$  and the constant shifts  $h_j \in \mathbb{R}^d$  are parameters. Notice that the quantity above is  $d!$  times the (signed)  $d$ -volume of the simplex spanned by  $(u_0, \dots, u_d)$ . Thus the estimate below (where of course  $\rho_j(t, y) = \rho(t, y + h_j)$ ) is useful only when  $(h_0, \dots, h_d)$  are affinely independent

**Theorem 1.3** *Consider an admissible inviscid gas flow in  $Q_T$ , with finite total mass and energy. Then one has*

$$(3) \quad \sup_{h_0, \dots, h_d} \int_{Q_T} \left( \left( \prod_{j=0}^d \rho_j \right) \cdot (\text{Cor}(u_0, \dots, u_d))^2 \right)^{\frac{1}{d}}(t, y) dy dt \leq_d M^{\frac{1}{d}} \sqrt{ME_0}.$$

We point out that the integrand in (3) shares the same physical dimension as the expected  $\rho^{1+\frac{1}{d}}|u|^2$ . Recall in passing that the product  $ME_0$  in the right-hand side of (3) may be replaced by its Galilean invariant version, exactly like in [6]:

$$\frac{1}{4} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \rho(0, y) \rho(0, z) |u(0, z) - u(0, y)|^2 dy dz + M \int_{\mathbb{R}^d} (\rho e)(0, y) dy,$$

where the last integral, the internal energy, remains unchanged.

Once again (3) has the flavor of a Strichartz inequality: denoting the space integral

$$H(t, h_0, \dots, h_d) := \int_{\mathbb{R}^d} \left( \left( \prod_{j=0}^d \rho_j \right) \cdot (\text{Cor}(u_0, \dots, u_d))^2 \right)^{\frac{1}{d}}(t, y) dy,$$

it tells us that  $H \in L_{h_0, \dots, h_d}^\infty L_t^1$ . This must be put in front of a direct application of (2), which gives instead  $H \in L_{t, h_1, \dots, h_d}^\infty L_{h_0}^d$ , see Paragraph 4.2.

We observe, sadly, that our collection of *a priori* estimates, namely the conservation of mass, decay of energy, together with (1) and (3), is not strong enough to give a meaning to the flux in the balance law of energy (which is a conservation law in some cases):

$$(4) \quad \partial_t \left( \frac{1}{2} \rho |u|^2 + \rho e \right) + \text{div}_y \left[ \left( \frac{1}{2} \rho |u|^2 + \rho e + p \right) u \right] \leq 0.$$

As a matter of fact, we control quantities which either are quadratic in the velocity field, or do not involve it, while the flux in (4) is cubic in  $u$ .

Another significant improvement of the corpus of estimates is the following statement concerning the pressure.

**Theorem 1.4** *Admissible flows of Gas dynamics, with finite total mass and energy, satisfy*

$$(5) \quad \sup_{\tau, \eta} \int_0^\infty \int_{\mathbb{R}^d} \left( \frac{(t - \tau)^2}{(E_0(t - \tau)^2 + M|y - \eta|^2)^{\frac{d}{2} + 1}} \right)^{\frac{1}{d}} p(t, y) dy dt \leq_d E_0^{1 - \frac{1}{d}}.$$

Inequality (5) is reminiscent of a well-known fact about convolution of functions  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  of bounded variations, say with compact support,

$$(6) \quad \left\| \frac{1}{|x|} * f \right\|_{L^\infty} \leq_n TV(f).$$

**Remark 1.1** *According to L. Tartar (personal communication), the latter is a consequence of the improved Sobolev–Lorentz embedding (see A. Alvino [1])*

$$(7) \quad BV(\mathbb{R}^n) \subset L^{\frac{n}{n-1}, 1}(\mathbb{R}^n),$$

with the fact that  $x \mapsto \frac{1}{|x|}$  belongs to the dual space  $L^{n, \infty}(\mathbb{R}^n)$ . Luc pointed out that the convolution product actually belongs to  $C_b(\mathbb{R}^n)$ , the space of continuous bounded functions, because of the density of  $C_K(\mathbb{R}^n)$  in  $L^{\frac{n}{n-1}, 1}(\mathbb{R}^n)$ .

We shall give below a new proof of (6), by means of Compensated Integrability, and extend it to the convolution by arbitrary positively homogeneous kernels of degree  $-1$ , see Proposition 3.1.

**Plan of the paper.** Section 2 displays new forms of Compensated Integrability. The homogeneous polynomial  $A \mapsto \det A$  being hyperbolic over  $\mathbf{Sym}_n^+$  in the sense of Gårding [3], we may use the properties of its multi-linearization, the *mixed determinant*. Part 3, which applies the abstract results to scalar functions, is two-fold. It begins with BV-functions and continues with a Gagliardo-like inequality for time-dependent functions. Section 4 treats applications to Gas dynamics, and contains the proofs of Theorems 1.3 and 1.4. In the same spirit, it displays a new estimate of the so-called *defect measure* introduced by [5] when averaging renormalized solutions of Boltzman equation.

**Acknowledgement.** I am grateful to Luc Tartar for sharing his precious knowledge of the history of Sobolev–Lorentz embeddings, and to P. Mironescu for his interest in this work.

## 2 New forms of Compensated Integrability

We start with a refined result, whose proof illustrates the scaling technique which is at stake almost everywhere in the theory.

**Theorem 2.1** *Let  $A \succ 0_d$  be Div-BV over  $\mathbb{R}^n$ . Then*

$$(8) \quad \int_{\mathbb{R}^n} (\det A)^{\frac{1}{n-1}} dx \leq n^{\frac{n}{n-1}} c_n \left( \prod_{i=1}^n \|(\operatorname{Div} A)_i\|_{\mathcal{M}} \right)^{\frac{1}{n-1}}.$$

*Proof*

We rescale both dependent and independent variables,

$$x'_j = \mu_j x_j, \quad a'_{ij} = \mu_i \mu_j a_{ij}.$$

Since  $A' = \Delta A \Delta$  for a diagonal matrix  $\Delta$ , it is still positive semi-definite. Because  $\partial'_j = \frac{1}{\mu_j} \partial_j$ , we have

$$(\operatorname{Div}' A')_i = \mu_i (\operatorname{Div} A)_i$$

and thus  $A'$  is also Div-BV. Let us apply Thm 1.1 to  $A'$ , using

$$\det A' = \left( \prod_{j=1}^n \mu_j \right)^2 \det A, \quad dx' = \left( \prod_{j=1}^n \mu_j \right) dx,$$

and

$$|\operatorname{Div} A| \leq \sum_{j=1}^n |(\operatorname{Div} A)_j|.$$

We obtain

$$\int_{\mathbb{R}^n} (\det A)^{\frac{1}{n-1}} dx \leq c_n \left( \prod_{j=1}^n \mu_j \right)^{-\frac{1}{n-1}} \left( \sum_{i=1}^n \mu_i \|(\operatorname{Div} A)_i\|_{\mathcal{M}} \right)^{\frac{n}{n-1}}.$$

Choosing

$$\mu_i = \frac{1}{\|(\operatorname{Div} A)_i\|_{\mathcal{M}}},$$

we obtain (8). ■

**Remark 2.1** *The inequality in Theorem 2.1 can be recast as*

$$\log \int_{\mathbb{R}^n} (\det A)^{\frac{1}{n-1}} dx \leq \log(n^{\frac{n}{n-1}} c_n) + \frac{1}{n-1} \sum_{j=1}^n \log \|(\operatorname{Div} A)_j\|_{\mathcal{M}}.$$

*Since it remains valid when the canonical basis is replaced by an arbitrary orthonormal basis, we obtain by averaging*

$$\log \int_{\mathbb{R}^n} (\det A)^{\frac{1}{n-1}} dx \leq \log(n^{\frac{n}{n-1}} c_n) + \frac{n}{n-1} \int_{S_{n-1}} \log \|(\operatorname{Div} A) \cdot e\|_{\mathcal{M}} ds(e).$$

*In other words, we have*

$$(9) \quad \int_{\mathbb{R}^n} (\det A)^{\frac{1}{n-1}} dx \leq c_n \left( n \exp \int_{S_{n-1}} \log \|\operatorname{div}(Ae)\|_{\mathcal{M}} ds(e) \right)^{\frac{n}{n-1}}.$$

## 2.1 Estimating mixed determinant.

To go further, we invoke the notion of *mixed determinant*. This is the symmetric  $n$ -linear form  $D_n$  with the property that

$$D_n(M, \dots, M) = \det M, \quad \forall M \in \mathbf{M}_n(\mathbb{R}).$$

For instance, the case  $n = 2$  (where the determinant is a quadratic form) gives

$$D_2(M, M') = \frac{1}{2} (\det(M + M') - \det M - \det M') = \frac{1}{4} (\det(M + M') - \det(M - M')).$$

More generally, we have

$$D_n(M_1, \dots, M_n) = \frac{1}{2^n n!} \sum_{\varepsilon_1, \dots, \varepsilon_n = \pm 1} \varepsilon_1 \cdots \varepsilon_n \det(\varepsilon_1 M_1 + \cdots + \varepsilon_n M_n).$$

We notice the useful instance

$$(10) \quad D_n(B, M, \dots, M) = \frac{1}{n} \operatorname{Tr}(B^T \widehat{M}).$$



**Hyperbolicity.** Let us recall that the homogeneous polynomial  $\det : \mathbf{Sym}_n \rightarrow \mathbb{R}$  is *hyperbolic* in Gårding's terminology [3], meaning that there exists a direction  $E \neq 0_n$  such that  $\det E > 0$  and for every  $A \in \mathbf{Sym}_n$ , the roots of the univariate polynomial  $t \mapsto \det(A - tE)$  are real. The connected component  $\Gamma$  of  $E$  in the complement of  $\{A \mid \det A = 0\}$  is a convex cone, called the *forward cone*. Actually, the hyperbolic property stands in every direction  $E' \in \Gamma$ . We may of course choose  $E = I_n$ , and we see that  $\Gamma = \mathbf{SPD}_n$  is the positive definite cone.

Gårding proved that a hyperbolic polynomial satisfies the reverse Hölder inequality in the closure of its forward cone. In the case of the determinant, this reads

**Proposition 2.1** *For every  $A_1, \dots, A_n \in \mathbf{Sym}_n^+$ , there holds*

$$(11) \quad \left( \prod_{j=1}^n \det A_j \right)^{\frac{1}{n}} \leq D_n(A_1, \dots, A_n).$$

In particular the mixed determinant takes non-negative values over  $\mathbf{Sym}_n^+ \times \dots \times \mathbf{Sym}_n^+$ . Developing the multilinear expression  $D_n(A_1 + \dots + A_n, \dots, A_1 + \dots + A_n)$ , we deduce

$$(12) \quad D_n(A_1, \dots, A_n) \leq \frac{1}{n!} \det(A_1 + \dots + A_n), \quad \forall A_1, \dots, A_n \in \mathbf{Sym}_n^+.$$

Applying Theorem 1.1 to  $A = A_1 + \dots + A_n$ , and using (12), we deduce an inequality for non-negative Div-BV tensors over  $\mathbb{R}^n$ :

$$\int_{\mathbb{R}^n} (D_n(A_1, \dots, A_n))^{\frac{1}{n-1}} dx \leq_n \left( \sum_{j=1}^n \|\mathrm{Div} A_j\|_{\mathcal{M}} \right)^{\frac{n}{n-1}}.$$

Replacing  $A_j$  by  $\lambda_j A_j$  where  $\lambda_j = \|\mathrm{Div} A_j\|_{\mathcal{M}}^{-1}$ , we infer the estimate

**Theorem 2.2** *Given positive semi-definite Div-BV tensors  $A_1, \dots, A_n$  over  $\mathbb{R}^n$ , one has*

$$(13) \quad \int_{\mathbb{R}^n} (D_n(A_1, \dots, A_n))^{\frac{1}{n-1}} dx \leq_n \left( \prod_{j=1}^n \|\mathrm{Div} A_j\|_{\mathcal{M}} \right)^{\frac{1}{n-1}}.$$

## 2.2 Estimating Schur complements

Consider a positive semi-definite Div-BV tensor written blockwise

$$A = \begin{pmatrix} \rho & m^T \\ m & B \end{pmatrix},$$

where  $\rho$  is scalar. The positiveness tells us on the one hand that  $\rho \geq 0$  and on the other hand that either  $\rho = 0$ ,  $m = 0$  and  $B \succ 0_{n-1}$ , or  $\rho > 0$  and  $B \succ \frac{m \otimes m}{\rho}$ . Thus let us assume that  $\rho > 0$ , and denote  $S := B - \frac{m \otimes m}{\rho} \succ 0_{n-1}$  its *Schur complement*. Then  $A$  splits into the sum of two positive semi-definite matrices:

$$(14) \quad A = \begin{pmatrix} \rho & m^T \\ m & \frac{m \otimes m}{\rho} \end{pmatrix} + \begin{pmatrix} 0 & 0^T \\ 0 & S \end{pmatrix} =: \begin{pmatrix} \rho & m^T \\ m & \frac{m \otimes m}{\rho} \end{pmatrix} + A'.$$

The *Schur complement formula*

$$(15) \quad \det A = \rho \det S$$

allows us to make calculations without expressing  $S$  in closed form.

Suppose now that  $F \succ 0_n$  is another Div-BV tensor. Developing the expression  $D_n(F, A, \dots, A)$  and discarding all non-negative terms but one, we obtain

$$(16) \quad D_n(F, A, \dots, A) \geq D_n(F, A', \dots, A') = \frac{1}{n} \text{Tr}(F \widehat{A}') = \frac{1}{n} f_{11} \det S.$$

We infer the Schur complement estimate:

**Proposition 2.2** *Given two positive semi-definite Div-BV tensors  $F, A$  over  $\mathbb{R}^n$ , with  $a_{11} > 0$ , the Schur complement  $S$  of  $a_{11}$  satisfies*

$$(17) \quad \int_{\mathbb{R}^n} (f_{11} \det S)^{\frac{1}{n-1}} dx \leq_n \|\text{Div} A\|_{\mathcal{M}} \|\text{Div} F\|_{\mathcal{M}}^{\frac{1}{n-1}}.$$

**Remark 2.2** *When the entry  $a_{11}$  is not strictly positive everywhere, it might sometimes be difficult, if not impossible, to define its Schur complement. Instead, it may happen that the tensor decomposes as  $A = K + A'$  with  $K \succ 0_n$  and*

$$A' = \begin{pmatrix} 0 & 0 \\ 0 & \Sigma \end{pmatrix}, \quad \Sigma \succ 0_{n-1}.$$

*Then the same argument as above works: (16) still holds true, and implies a generalization of (17),*

$$(18) \quad \int_{\mathbb{R}^n} (f_{11} \det \Sigma)^{\frac{1}{n-1}} dx \leq_n \|\text{Div} A\|_{\mathcal{M}} \|\text{Div} F\|_{\mathcal{M}}^{\frac{1}{n-1}}.$$

The full strength of (18) occurs when we choose an extreme tensor  $F$ . By “extreme”, we mean that  $F$  be positively homogeneous of degree  $1 - n$ , see [7]. Since the homogeneity is not compatible with  $f_{ij} \in L^1(\mathbb{R}^n)$ , we use a cut-off function  $\phi(r)$  before getting rid of it:

$$F = \phi(r)G, \quad G(x) = \frac{x \otimes x}{r^{n+1}}, \quad r = |x|.$$

The cut-off is a non-increasing function such that  $\phi(r) \equiv 1$  over  $(0, R)$  and  $\equiv 0$  over  $(R + 1, +\infty)$ . Since  $\text{Div} G \equiv 0$ , we have

$$\|\text{Div} F\|_{\mathcal{M}} = \int_{\mathbb{R}^d} |G \nabla \phi| dx = \int_{\mathbb{R}^d} |\phi'(r)| \frac{dx}{r^{n-1}} = |S_{n-1}| \int_0^\infty |\phi'(r)| dr = |S_{n-1}|.$$

Applying (17), we obtain

$$\int_{|x| < R} \left( \frac{x_1^2}{r^{n+1}} \right)^{\frac{1}{n-1}} (\det \Sigma)^{\frac{1}{n-1}} dx \leq_n \|\text{Div} A\|_{\mathcal{M}}.$$

Taking the limit as  $R \rightarrow +\infty$  and relaxing the position of the origin, we obtain the estimate

$$(19) \quad \sup_{\xi \in \mathbb{R}^n} \int_{\mathbb{R}^n} \left( \frac{(x_1 - \xi_1)^2}{|x - \xi|^{n+1}} \right)^{\frac{1}{n-1}} (\det \Sigma)^{\frac{1}{n-1}} dx \leq_n \|\text{Div} A\|_{\mathcal{M}}.$$

When  $A$  is positive definite, the Schur complement is well-defined. Then applying a rotation, using (15) and the Schur formula, we conclude

**Theorem 2.3** *Let  $A : \mathbb{R}^n \rightarrow \mathbf{SPD}_n$  be a Div-BV tensor. Then we have*

$$(20) \quad \sup_{\omega \in S_{n-1}} \sup_{\xi \in \mathbb{R}^n} \int_{\mathbb{R}^n} \left( \frac{(\omega \cdot (x - \xi))^2}{|x - \xi|^{n+1}} \right)^{\frac{1}{n-1}} \left( \frac{\det A}{\omega^T A \omega} \right)^{\frac{1}{n-1}} dx \leq_n \|\text{Div} A\|_{\mathcal{M}}.$$

### 2.3 Estimating the rank-one part in (14)

Let us rewrite (14) in the form  $A = \rho U \otimes U + A'$ , where

$$U = \begin{pmatrix} 1 \\ u \end{pmatrix}, \quad u = \frac{m}{\rho}.$$

Suppose that we are given a collection  $(A_1, \dots, A_n)$  of positive semi-definite Div-BV tensors over  $\mathbb{R}^n$ . Let us form the, still positive and Div-BV, tensor

$$\mathbb{A} = A_1 + \dots + A_n.$$

Decomposing  $A_j = \rho_j U_j \otimes U_j + A'_j$ , we have

$$\mathbb{A} \succ \rho_1 U_1 \otimes U_1 + \cdots + \rho_n U_n \otimes U_n,$$

whence

$$\det \mathbb{A} \geq \det(\rho_1 U_1 \otimes U_1 + \cdots + \rho_n U_n \otimes U_n).$$

Expressing

$$\rho_1 U_1 \otimes U_1 + \cdots + \rho_n U_n \otimes U_n = U^T R U, \quad R = \text{diag}(\rho_1, \dots, \rho_n),$$

where  $U$  stands for the matrix whose columns are  $U_1, \dots, U_n$ , we infer

$$\det \mathbb{A} \geq (\det R) \cdot (\det U)^2 = \left( \prod_{j=1}^n \rho_j \right) \cdot [\text{Cor}(u_1, \dots, u_n)]^2.$$

Assembling the material above, and applying a standard scaling argument, we end up with

**Proposition 2.3** *Let  $A_1, \dots, A_n$  be positive semi-definite Div-BV tensors over  $\mathbb{R}^n$ , written block-wise*

$$A_j = \rho_j \begin{pmatrix} 1 \\ u_j \end{pmatrix} \otimes \begin{pmatrix} 1 \\ u_j \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & B_j \end{pmatrix}.$$

Then

$$\int_{\mathbb{R}^n} \left( \prod_{j=1}^n \rho_j \cdot [\text{Cor}(u_1, \dots, u_n)]^2 \right)^{\frac{1}{n-1}} dx \leq_n \left( \sum_{j=1}^n \|\text{Div} A_j\|_{\mathcal{M}} \right)^{\frac{n}{n-1}}.$$

Remark that the integrand in the left-hand side has the dimension of a  $\rho^{\frac{n}{n-1}} |u|^2$ .

## 3 Applications to scalar functions

### 3.1 BV functions

If  $f \in BV(\mathbb{R}^n)$  is non-negative, we may apply (13) to the choices

$$A_1 = F, \quad A_2 = \cdots = A_n = f I_n.$$

With (10), we find

$$D_n(F, f I_n, \dots, f I_n) = \frac{1}{n} f^{n-1} \text{Tr} F = f^{n-1} \frac{\phi}{nr^{n-1}}.$$

With  $\text{Div}(fI_n) = \nabla f$ , we thus obtain

$$\int_{\mathbb{R}^n} f(x) \phi(r) \frac{1}{r^{n-1}} \frac{dx}{r} \leq_n \|\text{Div}(fI_n)\|_{\mathcal{M}} = TV(f).$$

Letting  $R \rightarrow +\infty$ , this yields

$$\int_{\mathbb{R}^n} f(x) \frac{dx}{r} \leq_n TV(f).$$

Eventually, placing the origin at an arbitrary point, this rewrites

$$\forall \xi \in \mathbb{R}^n, \quad \int_{\mathbb{R}^n} f(x) \frac{dx}{|x - \xi|} \leq_n TV(f).$$

To pass from non-negative functions to signed functions, we may use  $|r^{-1} * f| \leq r^{-1} * |f|$ , and  $TV(|f|) \leq TV(f)$ . This re-proves (6).

The calculation above can be generalized as follows. If  $g \in L^{n-1}(S_{n-1})$  is non-negative, then choose instead

$$\bar{G}(x) = g\left(\frac{x}{r}\right)^{n-1} \frac{x \otimes x}{r^{n+1}}$$

and  $F(x) = \phi(r)\bar{G}(x)$  as before. In [7] we have seen that  $\text{Div } \bar{G} = V\delta_{x=0}$  where

$$V = \int_{S_{n-1}} g(\omega)^{n-1} \omega ds(\omega).$$

With  $\phi$  as before, this yields

$$\text{Div } F = \phi(0)V\delta_{x=0} + g\left(\frac{x}{r}\right)^{n-1} \frac{\phi'}{r^{n-1}} \vec{e}_r,$$

Whence

$$\|\text{Div } F\|_{\mathcal{M}} \leq 2\|g\|_{L^{n-1}(S_{n-1})}^{n-1}.$$

Then (13), applied to  $(F, fI_n, \dots, fI_n)$  yields

$$\int_{\mathbb{R}^n} f(x) \phi(r) \frac{1}{r^{n-1}} g\left(\frac{x}{r}\right) \frac{dx}{r} \leq_n TV(f) \|g\|_{L^{n-1}(S_{n-1})}.$$

Taking as above the supremum over the admissible  $\phi$  (the limit as  $R \rightarrow +\infty$ ), and choosing arbitrarily the origin, we end up with (see also Remark 1.1 for the replacement of  $L^\infty$  by  $C_b$ )

**Proposition 3.1** *There exists a finite constant  $C_n$  such that, for every  $g \in L^{n-1}(S_{n-1})$ , there holds*

$$\|\bar{g} * f\|_{L^\infty} \leq C_n \cdot TV(f) \|g\|_{L^{n-1}(S_{n-1})}, \quad \forall f \in BV(\mathbb{R}^n), \quad \bar{g}(x) := \frac{1}{r} g\left(\frac{x}{r}\right).$$

### 3.2 A Gagliardo-like inequality

Let us recall Gagliardo's inequality [4] in  $\mathbb{R}^d$ : given  $d$  functions  $f_j \in L^{d-1}(\mathbb{R}^{d-1})$ , the new function

$$f(y) = \prod_{j=1}^d f_j(\hat{y}_j), \quad y \in \mathbb{R}^d$$

is integrable. And there is a functional inequality

$$\|f\|_{L^1(\mathbb{R}^d)} \leq \prod_{j=1}^d \|f_j\|_{L^{d-1}(\mathbb{R}^{d-1})}.$$

If  $d = 2$ , this is nothing but Fubini's theorem. For general  $d$ , this can be recovered, up to a multiplicative constant, by applying (8) to the tensor  $\phi \text{diag}(|f_1|^{d-1}, \dots, |f_d|^{d-1})$ , where  $\phi$  is a cut-off as above, see [6].

We now introduce a time parameter, thus having  $d$  functions  $f_j(t, \hat{y}_j)$  and defining as above

$$(21) \quad f(t, y) = \prod_{j=1}^d f_j(t, \hat{y}_j), \quad (t, y) \in \mathbb{R}^{1+d}.$$

Mind that each  $f_j$  is defined over a  $d$ -dimensional space. Let us set  $n = 1 + d$  and form the tensor

$$A(x) = \frac{\phi(r)x \otimes x}{r^{n+1}} + \text{diag}(0, \phi(|y_1|)|f_1(\hat{x}_1)|^{n-1}, \dots, \phi(|y_d|)f_d(\hat{x}_d)^{n-1}) =: \sum_{i=0}^d A_i.$$

We have as above

$$\|\text{Div} A_0\|_{\mathcal{M}} = |S_{n-1}|,$$

while

$$\|\text{Div} A_j\|_{\mathcal{M}} = \int_{\mathbb{R}^n} |\phi'(|y_j|)| \cdot |f_j(\hat{x}_j)|^{n-1} dx = 2 \|f_j\|_{L^d(\mathbb{R}^d)}^d.$$

By Schur formula,

$$\det A = \frac{\phi(r)t^2}{r^{n+1}} \cdot \prod_{j=1}^d \phi(|y_j|)|f_j(\hat{x}_j)|^{n-1}.$$

Applying Theorem 1.1, we infer

$$\int_{|x| < R} \left( \frac{t^2}{(t^2 + |y|^2)^{1+\frac{d}{2}}} \right)^{\frac{1}{d}} \left| \prod_{j=1}^d f_j(t, \hat{y}_j) \right| dy dt \leq d \left( 1 + \sum_{j=1}^d \|f_j\|_{L^d(\mathbb{R} \times \mathbb{R}^{d-1})}^d \right)^{1+\frac{1}{d}}.$$

Passing to the limit as  $R \rightarrow +\infty$ , and making the appropriate scaling, we conclude

**Theorem 3.1** *Given the functions  $f_1, \dots, f_d \in L^d(\mathbb{R} \times \mathbb{R}^{d-1})$ , the function  $f$  defined over  $\mathbb{R} \times \mathbb{R}^d$  by (21) satisfies*

$$\int_{\mathbb{R} \times \mathbb{R}^d} \left( \frac{t^2}{(t^2 + |y|^2)^{1+\frac{d}{2}}} \right)^{\frac{1}{d}} |f(t,y)| dy dt \leq_d \prod_{j=1}^d \|f_j\|_{L^d(\mathbb{R} \times \mathbb{R}^{d-1})}.$$

## 4 Applications to Gas dynamics

Once again we set  $n = 1 + d$ , where  $d$  is the space dimension, so that  $\frac{1}{n-1} = \frac{1}{d}$ . The Euler system governs the conservation of mass and linear momentum

$$\begin{aligned} \partial_t \rho + \operatorname{div}_y(\rho u) &= 0, \\ \partial_t(\rho u) + \operatorname{Div}_y(\rho u \otimes u) + \nabla_y p &= 0. \end{aligned}$$

This is recast as  $\operatorname{Div}_{t,y} A = 0$ , where

$$A = \begin{pmatrix} \rho & \rho u^T \\ \rho u & \rho u \otimes u + p I_d \end{pmatrix} = \rho U \otimes U + p J \quad \text{in } Q_T,$$

with

$$U := \begin{pmatrix} 1 \\ u \end{pmatrix}, \quad J = \begin{pmatrix} 0 & 0 \\ 0 & I_d \end{pmatrix}.$$

We recognize the Schur complement  $p I_d$  of  $\rho$  in  $A$ , whose determinant is just  $p^d$ .

### 4.1 A new estimate of the pressure

Applying (19), we have

$$(22) \quad \sup_{\tau, \eta} \int_{Q_T} \left( \frac{(t-\tau)^2}{((t-\tau)^2 + |y-\eta|^2)^{\frac{d}{2}+1}} \right)^{\frac{1}{d}} p(t,y) dy dt \leq_d M + \sqrt{ME_0}.$$

Since the bound does not depend upon the length of the time interval, this estimate is valid over  $Q_\infty$ .

The estimate above is not homogeneous from a Physical point of view. To prove Theorem 1.4, we apply a scaling technique as in [6]: for every parameter  $\mu > 0$ , the dependent/independent variables

$$t' = t, \quad y' = \mu y, \quad \rho' = \rho, \quad u' = \mu u, \quad p' = \mu^2 p$$

define an admissible flow, to which we may apply (22). Chosing  $\mu = \sqrt{M/E_0}$ , we obtain (5).

Notice that we may rewrite (5) in terms of the square root mean velocity  $\bar{u} := \sqrt{2E_0/M}$ ,

$$\sup_{\tau, \eta} \int_{Q_\tau} \left( \frac{(t-\tau)^2}{\left( (t-\tau)^2 + \frac{|y-\eta|^2}{\bar{u}^2} \right)^{\frac{d}{2}+1}} \right)^{\frac{1}{d}} p(t,y) dy dt \leq_d E_0.$$

## 4.2 The velocity field

To get more information about the velocity field than just what is given by the energy estimate, we apply Proposition 2.3 to the tensors  $A_0, \dots, A_d$  obtained from  $A$  by a space shift:

$$A_j(t,y) := A(t,y+h_j).$$

Hereabove  $h_0, \dots, h_d$  are constant vectors. Of course the  $A_j$ 's are Div-BV whenever  $A$  is so. We thus obtain the inequality

$$\int_{Q_\tau} \left( \prod_{j=0}^d \rho(t,y+h_j) \cdot (\text{Cor}(u(t,y+h_0), \dots, u(t,y+h_d)))^2 \right)^{\frac{1}{d}} dy dt \leq_d \left( M + \sqrt{ME_0} \right)^{1+\frac{1}{d}}.$$

After the same scaling procedure as above, this gives Theorem 1.3.

**Discussion.** The integral depends only upon  $(h_0, \dots, h_d)$  modulo translations; we may thus assume  $h_0 = 0$  without loss of generality. Estimate (3) is a statement about the expression

$$H(t, h_1, \dots, h_d) = \int_{\mathbb{R}^d} \left( \rho \prod_{j=1}^d \rho_j \cdot (\text{Cor}(u, u_1, \dots, u_d))^2 \right)^{\frac{1}{d}} dy.$$

Namely, it says that

$$(23) \quad H \in L_{h_1, \dots, h_d}^\infty L_t^1.$$

We shall compare (23) with what can be said of  $H$  by using only the conservation of total mass and the decay of total energy. On the one hand, (3) is established through Compensated Integrability and thus requires that the mass-momentum of the fluid be positive semi-definite. As such, it applies to an inviscid fluid, as well as to the conservation of mass/momentum satisfied



by the renormalized solutions of the Boltzman equation (see [5]), but it does not apply to a viscous fluid. On the other hand, the pure mass-energy estimate below is valid in a much more general context, since it does not assume the positiveness of the mass-momentum tensor.

For this direct estimate, we first notice that,  $U = (U_0 | \cdots | U_d)$  being the same matrix as above,

$$|\det U| = \left| \sum_{k=0}^d (-1)^k \det(\dots, u_{k-1}, u_{k+1}, \dots) \right| \leq \sum_{k=0}^d \prod_{j \neq k} |u_j|.$$

We infer a majorization  $H \leq_d \sum_{k=0}^d H_k$  where

$$H_k(t, h_1, \dots, h_d) = \int_{\mathbb{R}^d} \left( \prod_{j=0}^d \rho_j \cdot \prod_{j \neq k} |u_j|^2 \right)^{\frac{1}{d}} dy.$$

The treatment of  $H_k$  depends on whether  $k = d$  or  $k < d$ , although it yields the same bound. Let us begin with the latter case. Applying the Hölder inequality with exponents  $(d, \dots, d)$ , we have

$$\begin{aligned} H_k^d &\leq \left( \prod_{j \neq k, d} \int_{\mathbb{R}^d} \rho_j |u_j|^2 dy \right) \cdot \int_{\mathbb{R}^d} \rho_k(t, y) (\rho |u|^2)(t, y + h_d) dy \\ &\leq (2E_0)^{d-1} \int_{\mathbb{R}^d} \rho_k(t, y) (\rho |u|^2)(t, y + h_d) dy. \end{aligned}$$

Integrating in  $h_d$  and using Fubini, this gives

$$\int_{\mathbb{R}^d} H_k(t, h_1, \dots, h_d)^d dh_d \leq_d (2E_0)^d \int_{\mathbb{R}^d} \rho_k(t, y) dy = M(2E_0)^d.$$

If instead  $k = d$ , we have

$$\begin{aligned} H_d^d &\leq \left( \prod_{j=1}^{d-1} \int_{\mathbb{R}^d} \rho_j |u_j|^2 dy \right) \cdot \int_{\mathbb{R}^d} \rho(t, y + h_d) (\rho |u|^2)(t, y) dy \\ &\leq (2E_0)^{d-1} \int_{\mathbb{R}^d} \rho(t, y + h_d) (\rho |u|^2)(t, y) dy, \end{aligned}$$

and again

$$\int_{\mathbb{R}^d} H_d(t, h_1, \dots, h_d)^d dh_d \leq_d M E_0^d.$$

The **direct** mass-energy bound for  $H$  is thus

$$(24) \quad \sup_{t, h_1, \dots, h_{d-1}} \int_{\mathbb{R}^d} H(t, h_1, \dots, h_d)^d dh_d \leq_d M E_0^d.$$

Thus the mass-energy conservation (or decay) provides the qualitative result that

$$(25) \quad H \in L_{t,h_1,\dots,h_{d-1}}^\infty L_{h_d}^d.$$

Comparing with (23), this amounts to replacing  $L_{h_d}^\infty L_t^1$  by  $L_t^\infty L_{h_d}^d$ .

Somehow the inequality (24) is brutal, in the sense that it applies to every quantity  $\mathcal{H}$  of the form

$$\mathcal{H}(t,h_1,\dots,h_d) = \int_{\mathbb{R}^d} \left( \prod_{j=0}^d \rho_j \cdot F(u_0,\dots,u_d) \right)^{\frac{1}{d}} dy$$

where  $F$  is a homogeneous polynomial of global degree  $2d$ , quadratic in each argument.

### 4.3 Renormalized solutions of Boltzman equation

Let us consider the Boltzman equation

$$(26) \quad (\partial_t + v \cdot \nabla_y) f(t,y,v) = Q[f(t,y,\cdot)], \quad (t,y) \in Q_T = (0,T) \times \mathbb{R}^d, v \in \mathbb{R}^d.$$

The Cauchy problem consists in finding a solution which fits an initial data  $f(0,y,v) = f_0(y,v)$ . The existence of distributional solutions to this problem is not known, except in space dimension  $d = 1$ . When  $d \geq 1$ , R. DiPerna & P.-L. Lions [2] proved instead the existence of a weaker notion of solutions, called *renormalized*. We shall not give a precise definition of this notion, and we content ourselves to recall that it implies, at the macroscopic level, the conservation of mass and a weak form of the conservation of momentum, in the sense that

$$(27) \quad \begin{aligned} \partial_t \rho + \operatorname{div}_y m &= 0, & \partial_t m + \operatorname{Div}_y \left( \int_{\mathbb{R}^d} f v \otimes v dv + \Sigma \right) &= 0 \\ \rho(t,y) &:= \int_{\mathbb{R}^d} f dv, & m(t,y) &:= \int_{\mathbb{R}^d} f v dv, & \Sigma &\succ 0_d. \end{aligned}$$

The quantities  $\rho, m$  are the mass density and linear momentum. Compared to what is formally expected, the second equation above contains an additional term  $\Sigma$ , called the *defect measure*, which takes values in  $\mathbf{Sym}_d^+$ ; see [5]. Finally, it is known that the total mass

$$M = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} f_0(y,v) dv dy \equiv \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} f(t,y,v) dv dy$$

is a constant of the motion, and the total energy

$$E(t) := \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} f(t,y,v) |v|^2 dv dy + \frac{1}{2} \int d\operatorname{Tr} \Sigma(t,\cdot)$$

is a non-increasing function of time and satisfies

$$E(t) \leq E_0 := \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} f_0(y, v) |v|^2 dv dy.$$

The equations (27) can be recast by saying that the following tensor

$$A = \begin{pmatrix} \int_{\mathbb{R}^d} f dv & \int_{\mathbb{R}^d} f v^T dv \\ \int_{\mathbb{R}^d} f v dv & \int_{\mathbb{R}^d} f v \otimes v dv + \Sigma \end{pmatrix}$$

is Div-free. For finite total mass and energy, the extension of  $A$  by  $0_n$  away from  $Q_T = (0, T) \times \mathbb{R}^d$  is Div-BV.

As mentioned in Remark 2.2, it might be difficult to construct the Schur complement of the density, if it vanishes here and there. Instead, decomposing  $A = B + A'$  where

$$B(t, y) = \int_{\mathbb{R}^d} f(t, y, v) \begin{pmatrix} 1 \\ v \end{pmatrix} \otimes \begin{pmatrix} 1 \\ v \end{pmatrix} dv, \quad A' = \begin{pmatrix} 0 & 0 \\ 0 & \Sigma \end{pmatrix},$$

we may apply (18) to obtain

$$\sup_{\tau, \eta} \int_0^T \int_{\mathbb{R}^d} \left( \frac{(t - \tau)^2}{((t - \tau)^2 + |y - \eta|^2)^{1+d/2}} \right)^{\frac{1}{d}} d(\det \Sigma)^{\frac{1}{d}} \leq_d M + \sqrt{ME_0},$$

where we recall that  $(\det \Sigma)^{\frac{1}{d}}$  is a finite measure, satisfying

$$0 \leq (\det \Sigma)^{\frac{1}{d}} \leq \frac{1}{d} \text{Tr} \Sigma.$$

Letting  $T \rightarrow +\infty$  (as usual, the upper bound does not depend upon the length of the time interval), and applying the usual scaling trick, we conclude that the defect measure is constrained by

$$(28) \quad \sup_{\tau, \eta} \int_0^T \int_{\mathbb{R}^d} \left( \frac{(t - \tau)^2}{(E_0(t - \tau)^2 + M|y - \eta|^2)^{1+d/2}} \right)^{\frac{1}{d}} d(\det \Sigma)^{\frac{1}{d}} \leq_d E_0^{1-\frac{1}{d}}.$$

Inequality (28) tells us that  $(\det \Sigma)^{\frac{1}{d}}$  is not too singular. For instance it does not charge points (no Dirac mass);  $\Sigma$  itself might charge points, but its density at a Dirac mass must be a singular matrix.

## References

- [1] A. Alvino. Sulla diseguaglianza di Sobolev in spazi di Lorentz. *Boll. Un. Mat. Ital. A*, **14** (1977), pp 148–156.
- [2] R. DiPerna, P.-L. Lions. On the Cauchy problem for the Boltzmann equation: global existence and weak stability results. *Annals of Math.*, **130** (1990), pp 321–366.
- [3] L. Gårding. An inequality for hyperbolic polynomials. *J. Math. Mech.*, **8** (1959), pp 957–965.
- [4] E. Gagliardo. Proprietà di alcune di funzioni in più variabili. *Ricerche Mat.*, **7** (1958), pp 102–137.
- [5] P.-L. Lions, N. Masmoudi. From Boltzmann equation to the Navier–Stokes and Euler equations II. *Arch Rat. Mech. & Anal.*, **158** (2001), pp 195–211.
- [6] D. Serre. Divergence-free positive symmetric tensors and fluid dynamics. *Annales de l'Institut Henri Poincaré (analyse non linéaire)*, **35** (2018), pp 1209–1234.
- [7] D. Serre. Compensated integrability. Applications to the Vlasov–Poisson equation and other models of mathematical physics. *J. Math. Pures & Appl.*, **127** (2019), pp 67–88.